

Sandoz	Business use only	Page 1 of 19
1.3.1 spc-label-pl - common-spc - 6,350 (NL/H/0856/002/IB/030 – response on RMS comments, NL/H/0856/001)		20160113
PIPERACILLIN SODIUM+TAZOBACTAM SODIUM 2 G + 250 MG; 4 G + 500 MG POWDER FOR SOLUTION FOR INFUSION		721-5483.00 721-5484.00 721-5484.01

1.3.1.1 Summary of Product Characteristics

1. NAME OF THE MEDICINAL PRODUCT

[nationally completed name 2 g / 0.25 g] powder for solution for infusion.

[nationally completed name 4 g / 0.5 g] powder for solution for infusion.

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

[nationally completed name 2 g / 0.25 g]:

Each vial contains piperacillin (as sodium salt) equivalent to 2 g and tazobactam (as sodium salt) equivalent to 0.25 g.

Each vial of [nationally completed name] 2 g / 0.25 g contains 4.72 mmol (109 mg) of sodium.

After reconstitution with 10 ml diluent the solution contains piperacillin 200 mg/ml and tazobactam 25 mg/ml.

[nationally completed name 4 g / 0.5 g]:

Each bottle contains piperacillin (as sodium salt) equivalent to 4 g and tazobactam (as sodium salt) equivalent to 0.5 g.

Each bottle of [nationally completed name] 4g / 0.5 g contains 9.44 mmol (217 mg) of sodium.

After reconstitution with 20 ml diluent the solution contains piperacillin 200 mg/ml and tazobactam 25 mg/ml.

3. PHARMACEUTICAL FORM

Powder for solution for infusion.

A white to off-white powder.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

Sandoz	Business use only	Page 2 of 19
1.3.1 spc-label-pl - common-spc - 6,350 (NL/H/0856/002/IB/030 – response on RMS comments, NL/H/0856/001)		20160113
PIPERACILLIN SODIUM+TAZOBACTAM SODIUM 2 G + 250 MG; 4 G + 500 MG POWDER FOR SOLUTION FOR INFUSION		721-5483.00 721-5484.00 721-5484.01

[nationally completed name] is indicated for the following infections in adults and children over 2 years of age (see sections 4.2 and 5.1):

Adults and adolescents

- Severe pneumonia including hospital-acquired and ventilator-associated pneumonia
- Complicated urinary tract infections (including pyelonephritis)
- Complicated intra-abdominal infections
- Complicated skin and soft tissue infections (including diabetic foot infections)

Treatment of patients with bacteraemia that occurs in association with, or is suspected to be associated with, any of the infections listed above.

[nationally completed name] may be used in the management of neutropenic patients with fever suspected to be due to a bacterial infection.

Children 2 to 12 years of age

- Complicated intra-abdominal infections

[nationally completed name] may be used in the management of neutropenic children with fever suspected to be due to a bacterial infection.

Consideration should be given to official guidance on the appropriate use of antibacterial agents.

4.2 Posology and method of administration

Posology

The dose and frequency of [nationally completed name] depends on the severity and localisation of the infection and expected pathogens.

Adult and adolescent patients

Infections

The usual dose is 4 g piperacillin / 0.5 g tazobactam given every 8 hours.

For nosocomial pneumonia and bacterial infections in neutropenic patients, the recommended dose is 4 g piperacillin / 0.5 g tazobactam administered every 6 hours. This regimen may also be applicable to treat patients with other indicated infections when particularly severe.

Sandoz	Business use only	Page 3 of 19
1.3.1 spc-label-pl - common-spc - 6,350 (NL/H/0856/002/IB/030 – response on RMS comments, NL/H/0856/001)		20160113
PIPERACILLIN SODIUM+TAZOBACTAM SODIUM 2 G + 250 MG; 4 G + 500 MG POWDER FOR SOLUTION FOR INFUSION		721-5483.00 721-5484.00 721-5484.01

The following table summarises the treatment frequency and the recommended dose for adult and adolescent patients by indication or condition:

Treatment frequency	[nationally completed name] 4 g / 0.5 g
Every 6 hours	Severe pneumonia
	Neutropenic adults with fever suspected to be due to a bacterial infection.
Every 8 hours	Complicated urinary tract infections (including pyelonephritis)
	Complicated intra-abdominal infections
	Skin and soft tissue infections (including diabetic foot infections)

Renal impairment

The intravenous dose should be adjusted to the degree of actual renal impairment as follows (each patient must be monitored closely for signs of substance toxicity; medicinal product dose and interval should be adjusted accordingly):

Creatinine clearance (ml/min)	[nationally completed name] (recommended dose)
> 40	No dose adjustment necessary
20-40	Maximum dose suggested: 4 g / 0.5 g every 8 hours
< 20	Maximum dose suggested: 4 g / 0.5 g every 12 hours

For patients on haemodialysis, one additional dose of piperacillin / tazobactam 2 g / 0.25 g should be administered following each dialysis period, because haemodialysis removes 30%-50% of piperacillin in 4 hours.

Hepatic impairment

No dose adjustment is necessary (see section 5.2).

Dose in elderly patients

No dose adjustment is required for the elderly with normal renal function or creatinine clearance values above 40 ml/min.

Paediatric population (2-12 years of age)

Infections

The following table summarises the treatment frequency and the dose per body weight for paediatric patients 2-12 years of age by indication or condition:

Sandoz	Business use only	Page 4 of 19
1.3.1 spc-label-pl - common-spc - 6,350 (NL/H/0856/002/IB/030 – response on RMS comments, NL/H/0856/001)		20160113
PIPERACILLIN SODIUM+TAZOBACTAM SODIUM 2 G + 250 MG; 4 G + 500 MG POWDER FOR SOLUTION FOR INFUSION		721-5483.00 721-5484.00 721-5484.01

Dose per weight and treatment frequency	Indication / condition
80 mg Piperacillin / 10 mg Tazobactam per kg body weight / every 6 hours	Neutropenic children with fever suspected to be due to bacterial infections*
100 mg Piperacillin / 12.5 mg Tazobactam per kg body weight / every 8 hours	Complicated intra-abdominal infections*

* Not to exceed the maximum 4 g / 0.5 g per dose over 30 minutes.

Renal impairment

The intravenous dose should be adjusted to the degree of actual renal impairment as follows (each patient must be monitored closely for signs of substance toxicity; medicinal product dose and interval should be adjusted accordingly):

Creatinine clearance (ml/min)	[nationally completed name] (recommended dose)
> 50	No dose adjustment needed.
≤ 50	70 mg piperacillin / 8.75 mg tazobactam / kg every 8 hours.

For children on haemodialysis, one additional dose of 40 mg piperacillin / 5 mg tazobactam / kg should be administered following each dialysis period.

Use in children aged below 2 years

The safety and efficacy of [nationally completed name] in children 0-2 years of age has not been established.

No data from controlled clinical studies are available.

Treatment duration

The usual duration of treatment for most indications is in the range of 5-14 days. However, the duration of treatment should be guided by the severity of the infection, the pathogen(s) and the patient's clinical and bacteriological progress.

Route of administration

[nationally completed name 2 g/250 mg] is administered by intravenous infusion (over 30 minutes).

[nationally completed name 4 g/500 mg] is administered by intravenous infusion (over 30 minutes).

Sandoz	Business use only	Page 5 of 19
1.3.1 spc-label-pl - common-spc - 6,350 (NL/H/0856/002/IB/030 – response on RMS comments, NL/H/0856/001)		20160113
PIPERACILLIN SODIUM+TAZOBACTAM SODIUM 2 G + 250 MG; 4 G + 500 MG POWDER FOR SOLUTION FOR INFUSION		721-5483.00 721-5484.00 721-5484.01

For reconstitution instructions, see section 6.6.

4.3 Contraindications

Hypersensitivity to the active substances or to any other penicillin-antibacterial agent.

History of acute severe allergic reaction to any other beta-lactam active substances (e.g. cephalosporin, monobactam or carbapenem).

4.4 Special warnings and precautions for use

The selection of piperacillin / tazobactam to treat an individual patient should take into account the appropriateness of using a broad-spectrum semi-synthetic penicillin based on factors such as the severity of the infection and the prevalence of resistance to other suitable antibacterial agents.

Before initiating therapy with [nationally completed name], careful inquiry should be made concerning previous hypersensitivity reactions to penicillins, other beta-lactam agents (e.g. cephalosporin, monobactam or carbapenem) and other allergens. Serious and occasionally fatal hypersensitivity (anaphylactic/anaphylactoid [including shock]) reactions have been reported in patients receiving therapy with penicillins, including piperacillin / tazobactam. These reactions are more likely to occur in persons with a history of sensitivity to multiple allergens. Serious hypersensitivity reactions require the discontinuation of the antibiotic, and may require administration of epinephrine and other emergency measures.

Antibiotic-induced pseudomembranous colitis may be manifested by severe, persistent diarrhoea which may be life-threatening. The onset of pseudomembranous colitis symptoms may occur during or after antibacterial treatment. In these cases [nationally completed name] should be discontinued.

Therapy with [nationally completed name] may result in the emergence of resistant organisms, which might cause super-infections.

Bleeding manifestations have occurred in some patients receiving beta-lactam antibiotics. These reactions sometimes have been associated with abnormalities of coagulation tests, such as clotting time, platelet aggregation and prothrombin time, and are more likely to occur in patients with renal failure. If bleeding manifestations occur, the antibiotic should be discontinued and appropriate therapy instituted.

Sandoz	Business use only	Page 6 of 19
1.3.1 spc-label-pl - common-spc - 6,350 (NL/H/0856/002/IB/030 – response on RMS comments, NL/H/0856/001)		20160113
PIPERACILLIN SODIUM+TAZOBACTAM SODIUM 2 G + 250 MG; 4 G + 500 MG POWDER FOR SOLUTION FOR INFUSION		721-5483.00 721-5484.00 721-5484.01

Leukopenia and neutropenia may occur, especially during prolonged therapy; therefore, periodic assessment of haematopoietic function should be performed.

As with treatment with other penicillins, neurological complications in the form of convulsions may occur when high doses are administered, especially in patients with impaired renal function.

Each vial of [nationally completed name 2 g/250 mg] contains 4.72 mmol (109 mg) of sodium and each bottle of [nationally completed name 4 g/500 mg] contains 9.44 mmol (217 mg) of sodium. This should be taken into consideration for patients who are on a controlled sodium diet.

Hypokalaemia may occur in patients with low potassium reserves or those receiving concomitant medicinal products that may lower potassium levels; periodic electrolyte determinations may be advisable in such patients.

4.5 Interaction with other medicinal products and other forms of interaction

Non-depolarising muscle relaxants

Piperacillin when used concomitantly with vecuronium has been implicated in the prolongation of the neuromuscular blockade of vecuronium. Due to their similar mechanisms of action, it is expected that the neuromuscular blockade produced by any of the non-depolarising muscle relaxants could be prolonged in the presence of piperacillin.

Oral anticoagulants

During simultaneous administration of heparin, oral anticoagulants and other substances that may affect the blood coagulation system including thrombocyte function, appropriate coagulation tests should be performed more frequently and monitored regularly.

Methotrexate

Piperacillin may reduce the excretion of methotrexate; therefore, serum levels of methotrexate should be monitored in patients to avoid substance toxicity.

Probenecid

As with other penicillins, concurrent administration of probenecid and piperacillin / tazobactam produces a longer half-life and lower renal clearance for both piperacillin and tazobactam; however, peak plasma concentrations of either substances are unaffected.

Aminoglycosides

Piperacillin, either alone or with tazobactam, did not significantly alter the pharmacokinetics of tobramycin in subjects with normal renal function and with mild or moderate renal

Sandoz	Business use only	Page 7 of 19
1.3.1 spc-label-pl - common-spc - 6,350 (NL/H/0856/002/IB/030 – response on RMS comments, NL/H/0856/001)		20160113
PIPERACILLIN SODIUM+TAZOBACTAM SODIUM 2 G + 250 MG; 4 G + 500 MG POWDER FOR SOLUTION FOR INFUSION		721-5483.00 721-5484.00 721-5484.01

impairment. The pharmacokinetics of piperacillin, tazobactam, and the M1 metabolite were also not significantly altered by tobramycin administration.

The inactivation of tobramycin and gentamicin by piperacillin has been demonstrated in patients with severe renal impairment.

For information related to the administration of piperacillin / tazobactam with aminoglycosides please refer to section 6.2.

Vancomycin

No pharmacokinetic interactions have been noted between piperacillin / tazobactam and vancomycin.

Effects on laboratory tests

Non-enzymatic methods of measuring urinary glucose may lead to false-positive results, as with other penicillins. Therefore, enzymatic urinary glucose measurement is required under [nationally completed name] therapy.

A number of chemical urine protein measurement methods may lead to false-positive results. Protein measurement with dip sticks is not affected.

The direct Coombs test may be positive.

Bio-Rad Laboratories *Platelia Aspergillus* EIA tests may lead to false-positive results for patients receiving [nationally completed name]. Cross-reactions with non-*Aspergillus* polysaccharides and polyfuranoses with Bio-Rad Laboratories *Platelia Aspergillus* EIA test have been reported.

Positive test results for the assays listed above in patients receiving [nationally completed name] should be confirmed by other diagnostic methods.

4.6 Fertility, pregnancy and lactation

Pregnancy

There are no or a limited amount of data from the use of [nationally completed name] in pregnant women.

Studies in animals have shown developmental toxicity, but no evidence of teratogenicity, at doses that are maternally toxic (see section 5.3).

Sandoz	Business use only	Page 8 of 19
1.3.1 spc-label-pl - common-spc - 6,350 (NL/H/0856/002/IB/030 – response on RMS comments, NL/H/0856/001)		20160113
PIPERACILLIN SODIUM+TAZOBACTAM SODIUM 2 G + 250 MG; 4 G + 500 MG POWDER FOR SOLUTION FOR INFUSION		721-5483.00 721-5484.00 721-5484.01

Piperacillin and tazobactam cross the placenta. Piperacillin / tazobactam should only be used during pregnancy if clearly indicated, i.e. only if the expected benefit outweighs the possible risks to the pregnant woman and foetus.

Breast-feeding

Piperacillin is excreted in low concentrations in human milk; tazobactam concentrations in human milk have not been studied. Women who are breast feeding should be treated only if the expected benefit outweighs the possible risks to the woman and child.

Fertility

A fertility study in rats showed no effect on fertility and mating after intraperitoneal administration of tazobactam or the combination piperacillin / tazobactam (see section 5.3).

4.7 Effects on ability to drive and use machines

No studies on the effect on the ability to drive and use machines have been performed.

4.8 Undesirable effects

The most commonly reported adverse reactions (occurring in 1 to 10 patients in 100) are diarrhoea, vomiting, nausea and rash.

In the following table, adverse reactions are listed by system organ class and MedDRA-preferred term. Within each frequency grouping, undesirable effects are presented in order of decreasing seriousness.

System Organ Class	Common ≥ 1/100 to < 1/10	Uncommon ≥ 1/1,000 to < 1/100	Rare ≥ 1/10,000 to < 1/1,000	Very rare (< 1/10,000)
Infections and infestations		candidal superinfection		
Blood and lymphatic system disorders		leukopenia, neutropenia, thrombocytopenia	anaemia, haemolytic anaemia, purpura, epistaxis, bleeding time prolonged, eosinophilia	agranulocytosis, pancytopenia, activated partial thromboplastin time prolonged, prothrombin time prolonged, Coombs direct test positive,

Sandoz	Business use only	Page 9 of 19
1.3.1 spc-label-pl - common-spc - 6,350 (NL/H/0856/002/IB/030 – response on RMS comments, NL/H/0856/001)		20160113
PIPERACILLIN SODIUM+TAZOBACTAM SODIUM 2 G + 250 MG; 4 G + 500 MG POWDER FOR SOLUTION FOR INFUSION		721-5483.00 721-5484.00 721-5484.01

System Organ Class	Common ≥ 1/100 to < 1/10	Uncommon ≥ 1/1,000 to < 1/100	Rare ≥ 1/10,000 to < 1/1,000	Very rare (< 1/10,000)
				thrombocytopenia
Immune system disorders		hypersensitivity	anaphylactic/ anaphylactoid reaction (including shock)	
Metabolism and nutrition disorders				hypokalaemia, blood glucose decreased, blood albumin decreased, blood protein total decreased
Nervous system disorders		headache, insomnia		
Vascular disorders		hypotension, thrombophlebitis, phlebitis	flushing	
Gastrointestinal disorders	diarrhoea, vomiting, nausea	jaundice, stomatitis, constipation, dyspepsia	pseudo- membranous colitis, abdominal pain	
Hepatobiliary disorders		alanine aminotransferase increased, aspartate aminotransferase increased	hepatitis, blood bilirubin increased, blood alkaline phosphatase increased, gamma- glutamyltrans- ferase increased	
Skin and subcutaneous tissue disorders	rash, including maculopapular rash	urticaria, pruritus	erythema multiforme, dermatitis bullous, exanthema	toxic epidermal necrolysis, Stevens-Johnson syndrome
Musculoskeletal and connective tissue disorders			arthralgia, myalgia	
Renal and urinary disorders		blood creatinine increased	renal failure, tubulointerstitial nephritis	blood urea increased
General		pyrexia, injection-	chills	

Sandoz	Business use only	Page 10 of 19
1.3.1 spc-label-pl - common-spc - 6,350 (NL/H/0856/002/IB/030 – response on RMS comments, NL/H/0856/001)		20160113
PIPERACILLIN SODIUM+TAZOBACTAM SODIUM 2 G + 250 MG; 4 G + 500 MG POWDER FOR SOLUTION FOR INFUSION		721-5483.00 721-5484.00 721-5484.01

System Organ Class	Common ≥ 1/100 to < 1/10	Uncommon ≥ 1/1,000 to < 1/100	Rare ≥ 1/10,000 to < 1/1,000	Very rare (< 1/10,000)
disorders and administration site conditions		site reaction		

Piperacillin therapy has been associated with an increased incidence of fever and rash in cystic fibrosis patients.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorization of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system listed in Appendix V.

4.9 Overdose

Symptoms

There have been post-marketing reports of overdose with piperacillin / tazobactam. The majority of those events experienced, including nausea, vomiting and diarrhoea, have also been reported with the usual recommended dose. Patients may experience neuromuscular excitability or convulsions if higher than recommended doses are given intravenously (particularly in the presence of renal failure).

Treatment

In the event of an overdose, piperacillin / tazobactam treatment should be discontinued. No specific antidote is known.

Treatment should be supportive and symptomatic according to the patient's clinical presentation.

Excessive serum concentrations of either piperacillin or tazobactam may be reduced by haemodialysis (see section 4.4).

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Sandoz	Business use only	Page 11 of 19
1.3.1 spc-label-pl - common-spc - 6,350 (NL/H/0856/002/IB/030 – response on RMS comments, NL/H/0856/001)		20160113
PIPERACILLIN SODIUM+TAZOBACTAM SODIUM 2 G + 250 MG; 4 G + 500 MG POWDER FOR SOLUTION FOR INFUSION		721-5483.00 721-5484.00 721-5484.01

Pharmacotherapeutic group: Antibacterials for systemic use, Combinations of penicillins incl. beta-lactamase inhibitors; ATC code: J01C R05.

Mechanism of action

Piperacillin, a broad spectrum, semisynthetic penicillin exerts bactericidal activity by inhibition of both septum and cell wall synthesis.

Tazobactam, a beta-lactam structurally related to penicillins, is an inhibitor of many beta-lactamases, which commonly cause resistance to penicillins and cephalosporins, but it does not inhibit AmpC enzymes or metallo beta-lactamases. Tazobactam extends the antibiotic spectrum of piperacillin to include many beta-lactamase-producing bacteria that have acquired resistance to piperacillin alone.

Pharmacokinetic / Pharmacodynamic relationship

The time above the minimum inhibitory concentration (T>MIC) is considered to be the major pharmacodynamic determinant of efficacy for piperacillin.

Mechanism of resistance

The two main mechanisms of resistance to piperacillin / tazobactam are:

- Inactivation of the piperacillin component by those beta-lactamases that are not inhibited by tazobactam: beta-lactamases in the Molecular class B, C and D. In addition, tazobactam does not provide protection against extended-spectrum beta-lactamases (ESBLs) in the Molecular class A and D enzyme groups.
- Alteration of penicillin-binding proteins (PBPs), which results in the reduction of the affinity of piperacillin for the molecular target in bacteria.

Additionally, alterations in bacterial membrane permeability, as well as expression of multi-drug efflux pumps, may cause or contribute to bacterial resistance to piperacillin / tazobactam, especially in Gram-negative bacteria.

Breakpoints

EUCAST Clinical MIC Breakpoints for Piperacillin/Tazobactam (Version 2.0, 2012-01-01)

Pathogen	MIC breakpoint (mg/L)	
	S ≤	R >
Enterobacteriaceae	8 ¹	16 ¹
<i>Pseudomonas</i> spp. ²	16 ³	16 ³

Sandoz	Business use only	Page 12 of 19
1.3.1 spc-label-pl - common-spc - 6,350 (NL/H/0856/002/IB/030 – response on RMS comments, NL/H/0856/001)		20160113
PIPERACILLIN SODIUM+TAZOBACTAM SODIUM 2 G + 250 MG; 4 G + 500 MG POWDER FOR SOLUTION FOR INFUSION		721-5483.00 721-5484.00 721-5484.01

Pathogen	MIC breakpoint (mg/L)	
	S ≤	R >
Gram-positive anaerobes (except <i>Clostridium difficile</i>)	8 ¹	16 ¹
Gram-negative anaerobes	8 ¹	16 ¹
Non-species related breakpoints ⁴	4	16

¹For susceptibility testing purposes, the concentration of tazobactam is fixed at 4 mg/L.

²Breakpoints are based on high dose therapy (with or without tazobactam, 4 g x 4).

³For susceptibility testing purposes, the concentration of beta-lactamase inhibitor is fixed at 4 mg/L.

⁴Breakpoints apply to piperacillin-tazobactam dosage of 4 g x 3.

Susceptibility

The prevalence of acquired resistance may vary geographically and with time for selected species and local information on resistance is desirable, particularly when treating severe infections. As necessary, expert advice should be sought when the local prevalence of resistance is such that the utility of the agent in at least some types of infections is questionable.

Groupings of relevant species according to piperacillin / tazobactam susceptibility
COMMONLY SUSCEPTIBLE SPECIES
<u>Aerobic Gram-positive micro-organisms</u>
<i>Enterococcus faecalis</i>
<i>Listeria monocytogenes</i>
<i>Staphylococcus aureus</i> , methicillin-susceptible [‡]
<i>Staphylococcus</i> species, <i>coagulase negative</i> , methicillin-susceptible
<i>Streptococcus pyogenes</i>
<i>Group B streptococci</i>
<u>Aerobic Gram-negative micro-organisms</u>
<i>Citrobacter koseri</i>
<i>Haemophilus influenza</i>
<i>Moraxella catarrhalis</i>
<i>Proteus mirabilis</i>
<u>Anaerobic Gram-positive micro-organisms</u>

Sandoz	Business use only	Page 13 of 19
1.3.1 spc-label-pl - common-spc - 6,350 (NL/H/0856/002/IB/030 – response on RMS comments, NL/H/0856/001)		20160113
PIPERACILLIN SODIUM+TAZOBACTAM SODIUM 2 G + 250 MG; 4 G + 500 MG POWDER FOR SOLUTION FOR INFUSION		721-5483.00 721-5484.00 721-5484.01

<i>Clostridium</i> species
<i>Eubacterium</i> species
<i>Peptostreptococcus</i> species
<u>Anaerobic Gram-negative micro-organisms</u>
<i>Bacteroides fragilis</i> group
<i>Fusobacterium</i> species
<i>Porphyromonas</i> species
<i>Prevotella</i> species
SPECIES FOR WHICH ACQUIRED RESISTANCE MAY BE A PROBLEM
<u>Aerobic Gram-positive micro-organisms</u>
<i>Enterococcus faecium</i> ^{§,+}
<i>Streptococcus pneumonia</i>
<i>Streptococcus viridans</i> group
<u>Aerobic Gram-negative micro-organisms</u>
<i>Acinetobacter baumannii</i> [§]
<i>Burkholderia cepacia</i>
<i>Citrobacter freundii</i>
<i>Enterobacter</i> species
<i>Escherichia coli</i>
<i>Klebsiella pneumonia</i>
<i>Morganella morganii</i>
<i>Proteus vulgaris</i>
<i>Providencia</i> ssp.
<i>Pseudomonas aeruginosa</i>
<i>Serratia</i> species
INHERENTLY RESISTANT ORGANISMS
<u>Aerobic Gram-positive micro-organisms</u>
<i>Corynebacterium jeikeium</i>
<u>Aerobic Gram-negative micro-organisms</u>
<i>Legionella</i> species
<i>Stenotrophomonas maltophilia</i> ^{+,§}
<u>Other microorganisms</u>
<i>Chlamydia pneumonia</i>
<i>Mycoplasma pneumonia</i>

Sandoz	Business use only	Page 14 of 19
1.3.1 spc-label-pl - common-spc - 6,350 (NL/H/0856/002/IB/030 – response on RMS comments, NL/H/0856/001)		20160113
PIPERACILLIN SODIUM+TAZOBACTAM SODIUM 2 G + 250 MG; 4 G + 500 MG POWDER FOR SOLUTION FOR INFUSION		721-5483.00 721-5484.00 721-5484.01

[§] Species showing natural intermediate susceptibility.

⁺ Species for which high-resistance rates (more than 50%) have been observed in one or more areas/countries/regions within the EU.

[£] All methicillin-resistant staphylococci are resistant to piperacillin / tazobactam.

5.2 Pharmacokinetic properties

Absorption

The peak piperacillin and tazobactam concentrations after 4 g / 0.5 g administered over 30 minutes by intravenous infusion are 298 µg/ml and 34 µg/ml respectively.

Distribution

Both piperacillin and tazobactam are approximately 30% bound to plasma proteins. The protein binding of either piperacillin or tazobactam is unaffected by the presence of the other compound. Protein binding of the tazobactam metabolite is negligible.

Piperacillin / tazobactam is widely distributed in tissues and body fluids including intestinal mucosa, gallbladder, lung, bile, and bone. Mean tissue concentrations are generally 50 to 100% of those in plasma. Distribution into cerebrospinal fluid is low in subjects with non-inflamed meninges, as with other penicillins.

Biotransformation

Piperacillin is metabolised to a minor microbiologically active desethyl metabolite. Tazobactam is metabolised to a single metabolite that has been found to be micro-biologically inactive.

Elimination

Piperacillin and tazobactam are eliminated via the kidney by glomerular filtration and tubular secretion.

Piperacillin is excreted rapidly as unchanged substance, with 68% of the administered dose appearing in the urine. Tazobactam and its metabolite are eliminated primarily by renal excretion, with 80% of the administered dose appearing as unchanged substance and the remainder as the single metabolite. Piperacillin, tazobactam, and desethyl piperacillin are also secreted into the bile.

Sandoz	Business use only	Page 15 of 19
1.3.1 spc-label-pl - common-spc - 6,350 (NL/H/0856/002/IB/030 – response on RMS comments, NL/H/0856/001)		20160113
PIPERACILLIN SODIUM+TAZOBACTAM SODIUM 2 G + 250 MG; 4 G + 500 MG POWDER FOR SOLUTION FOR INFUSION		721-5483.00 721-5484.00 721-5484.01

Following single or multiple doses of piperacillin / tazobactam to healthy subjects, the plasma half-life of piperacillin and tazobactam ranged from 0.7 to 1.2 hours and was unaffected by dose or duration of infusion. The elimination half-lives of both piperacillin and tazobactam are increased with decreasing renal clearance.

There are no significant changes in piperacillin pharmacokinetics due to tazobactam. Piperacillin appears to slightly reduce the clearance of tazobactam.

Special populations

The half-life of piperacillin and of tazobactam increases by approximately 25% and 18%, respectively, in patients with hepatic cirrhosis compared to healthy subjects.

The half-life of piperacillin and tazobactam increases with decreasing creatinine clearance. The increase in half-life is two-fold and four-fold for piperacillin and tazobactam, respectively, at creatinine clearance below 20 ml/min compared to patients with normal renal function.

Haemodialysis removes 30% to 50% of piperacillin / tazobactam, with an additional 5% of the tazobactam dose removed as the tazobactam metabolite. Peritoneal dialysis removes approximately 6% and 21% of the piperacillin and tazobactam doses, respectively, with up to 18% of the tazobactam dose removed as the tazobactam metabolite.

Paediatric population

In a population PK analysis, estimated clearance for 9 month-old to 12 year-old patients was comparable to adults, with a population mean (SE) value of 5.64 (0.34) ml/min/kg. The piperacillin clearance estimate is 80% of this value for paediatric patients 2-9 months of age. The population mean (SE) for piperacillin volume of distribution is 0.243 (0.011) l/kg and is independent of age.

Elderly patients

The mean half-life for piperacillin and tazobactam were 32% and 55% longer, respectively, in the elderly compared with younger subjects. This difference may be due to age-related changes in creatinine clearance.

Race

No difference in piperacillin or tazobactam pharmacokinetics was observed between Asian (n=9) and Caucasian (n=9) healthy volunteers who received single 4 g / 0.5 g doses.

5.3 Preclinical safety data

Sandoz	Business use only	Page 16 of 19
1.3.1 spc-label-pl - common-spc - 6,350 (NL/H/0856/002/IB/030 – response on RMS comments, NL/H/0856/001)		20160113
PIPERACILLIN SODIUM+TAZOBACTAM SODIUM 2 G + 250 MG; 4 G + 500 MG POWDER FOR SOLUTION FOR INFUSION		721-5483.00 721-5484.00 721-5484.01

Preclinical data reveal no special hazard for humans based on conventional studies of repeated dose toxicity and genotoxicity. Carcinogenicity studies have not been conducted with piperacillin / tazobactam.

A fertility and general reproduction study in rats using intraperitoneal administration of tazobactam or the combination piperacillin / tazobactam reported a decrease in litter size and an increase in fetuses with ossification delays and variations of ribs, concurrent with maternal toxicity. Fertility of the F1 generation and embryonic development of F2 generation were not impaired.

Teratogenicity studies using intravenous administration of tazobactam or the combination piperacillin / tazobactam in mice and rats resulted in slight reductions in rat fetal weights at maternally toxic doses but did not show teratogenic effects.

Peri/postnatal development was impaired (reduced pup weights, increase in stillbirths, increase in pup mortality) concurrent with maternal toxicity after intraperitoneal administration of tazobactam or the combination piperacillin / tazobactam in the rat.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

None.

6.2 Incompatibilities

This medicinal product must not be mixed with other medicinal products except those mentioned in section 6.6.

This product must not be mixed or co-administrated with any aminoglycoside. The mixing of beta-lactam antibiotics with an aminoglycoside *in vitro* can result in substantial inactivation of the aminoglycoside.

Piperacillin / tazobactam should not be mixed with other substances in a syringe or infusion bottle since compatibility has not been established.

Piperacillin / tazobactam should be administered through an infusion set separately from any other drugs unless compatibility is proven.

Sandoz	Business use only	Page 17 of 19
1.3.1 spc-label-pl - common-spc - 6,350 (NL/H/0856/002/IB/030 – response on RMS comments, NL/H/0856/001)		20160113
PIPERACILLIN SODIUM+TAZOBACTAM SODIUM 2 G + 250 MG; 4 G + 500 MG POWDER FOR SOLUTION FOR INFUSION		721-5483.00 721-5484.00 721-5484.01

Due to chemical instability, piperacillin / tazobactam should not be used in solutions containing only sodium bicarbonate.

Lactated Ringer's (Hartmann's) solution is not compatible with piperacillin / tazobactam.

Piperacillin / tazobactam should not be added to blood products or albumin hydrolysates.

6.3 Shelf life

Unopened:

2 years

After reconstitution (and dilution):

Chemical and physical in-use stability has been demonstrated for 24 hours at 20-25°C and for 48 hours at 2-8°C.

From a microbiological point of view, once opened, the product should be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would normally not be longer than 24 hours at 2-8°C, unless reconstitution/dilution has taken place in controlled and validated aseptic conditions.

6.4 Special precautions for storage

This medicinal product does not require any special storage conditions.

For storage conditions of the reconstituted/diluted medicinal product, see section 6.3.

6.5 Nature and contents of container

[nationally completed name 2 g / 0.25 g]:

30 ml vial glass type III, with halogenated butyl rubber stopper and aluminium overseal with flip-off cap.

Pack sizes of 1, 5, 10, 12 and 50 vials.

[nationally completed name 4 g / 0.5 g]:

100 ml bottle glass type II, with halogenated butyl rubber stopper and aluminium overseal with flip-off cap.

Sandoz	Business use only	Page 18 of 19
1.3.1 spc-label-pl - common-spc - 6,350 (NL/H/0856/002/IB/030 – response on RMS comments, NL/H/0856/001)		20160113
PIPERACILLIN SODIUM+TAZOBACTAM SODIUM 2 G + 250 MG; 4 G + 500 MG POWDER FOR SOLUTION FOR INFUSION		721-5483.00 721-5484.00 721-5484.01

50 ml bottle glass type II, with halogenated butyl rubber stopper and aluminium overseal with flip-off cap.

50 ml Injection vial (type III), with halogenated butyl rubber stopper and aluminium overseal with flip-off cap.

Pack sizes of 1, 5, 10, 12 and 50 bottles.

Not all pack sizes may be marketed.

6.6 Special precautions for disposal and other handling

The reconstitution and dilution is to be made under aseptic conditions. The solution is to be inspected visually for particulate matter and discoloration prior to administration. The solution should only be used if the solution is clear and free from particles.

Intravenous use

Reconstitute each vial with the volume of solvent shown in the table below, using one of the compatible solvents for reconstitution. Swirl until dissolved. When swirled constantly, reconstitution generally occurs within 3 minutes (for details on handling, please see below).

Content of vial	Volume of solvent* to be added to vial
2 g/0.25 g (2 g piperacillin and 0.25 g tacobactam)	10 ml
4 g/0.5 g (4 g piperacillin and 0.5 g tacobactam)	20 ml

* Compatible solvents for reconstitution:

- water for injection;
- sodium chloride 9 mg/ml (0.9 %) solution in water for injection;
- glucose 50 mg/ml (5 %) solution in water for injection;
- glucose 50 mg/ml (5 %) solution in sodium chloride 9 mg/ml (0.9%) solution.

The reconstituted solutions should be withdrawn from the vial by syringe. When reconstituted as directed, the vial contents withdrawn by syringe will provide the labelled amount of piperacillin and tazobactam.

The reconstituted solutions may be further diluted to the desired volume (e.g. 50 ml to 150 ml) with one of the following compatible solvents:

- sodium chloride 9 mg/ml (0.9 %) solution in water for injection;
- glucose 50 mg/ml (5 %) solution in water for injection;

Sandoz	Business use only	Page 19 of 19
1.3.1 spc-label-pl - common-spc - 6,350 (NL/H/0856/002/IB/030 – response on RMS comments, NL/H/0856/001)		20160113
PIPERACILLIN SODIUM+TAZOBACTAM SODIUM 2 G + 250 MG; 4 G + 500 MG POWDER FOR SOLUTION FOR INFUSION		721-5483.00 721-5484.00 721-5484.01

- dextran (grade 40) 60 mg/ml (6%) solution in sodium chloride 9 mg/ml (0.9%) solution.

See section 6.2 for incompatibilities.

Any unused product or waste material should be disposed in accordance with local requirements.

For single use only. Discard any unused solution.

7. MARKETING AUTHORISATION HOLDER

[Name and address to be completed nationally]

8. MARKETING AUTHORISATION NUMBER(S)

[To be completed nationally]

9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

[To be completed nationally]

10. DATE OF REVISION OF THE TEXT

[To be completed nationally]