

## SUMMARY OF PRODUCT CHARACTERISTICS

### 1. NAME OF THE MEDICINAL PRODUCT

<[Invented name]> 50 mg film-coated tablets  
<[Invented name]> 200 mg film-coated tablets

### 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each tablet contains 50 mg voriconazole.

Each tablet contains 200 mg voriconazole.

Excipient with known effect: each tablet contains 58.08 mg lactose.

Excipient with known effect: each tablet contains 232.30 mg lactose.

For the full list of excipients, see section 6.1.

### 3. PHARMACEUTICAL FORM

Film-coated tablet

White to off- white, round film coated tablets with diameter approx. 7.1 mm.

White to off- white, oval film coated tablets with dimensions approx. 15.8 x 7.9 mm.

### 4. CLINICAL PARTICULARS

#### 4.1 Therapeutic indications

Voriconazole is a broad spectrum, triazole antifungal agent and is indicated in adults and children aged 2 years and above as follows:

Treatment of invasive aspergillosis.

Treatment of candidaemia in non-neutropenic patients.

Treatment of fluconazole-resistant serious invasive *Candida* infections (including *C. krusei*).

Treatment of serious fungal infections caused by *Scedosporium* spp. and *Fusarium* spp.

<[Invented name]> should be administered primarily to patients with progressive, possibly life-threatening infections.

Prophylaxis of invasive fungal infections in high risk allogeneic hematopoietic stem cell transplant (HSCT) recipients.

## 4.2 Posology and method of administration

### Posology

Electrolyte disturbances such as hypokalaemia, hypomagnesaemia and hypocalcaemia should be monitored and corrected, if necessary, prior to initiation and during voriconazole therapy (see section 4.4).

<[Invented name]> is also available as 200 mg film-coated tablets.

<[Invented name]> is also available as 50 mg film-coated tablets.

### Treatment

#### *Adults*

Therapy must be initiated with the specified loading dose regimen of either intravenous or oral voriconazole to achieve plasma concentrations on Day 1 that are close to steady state. On the basis of the high oral bioavailability (96 %; see section 5.2), switching between intravenous and oral administration is appropriate when clinically indicated.

Detailed information on dosage recommendations is provided in the following table:

	Intravenous	Oral	
		Patients 40 kg and above*	Patients less than 40 kg*
<b>Loading dose regimen (first 24 hours)</b>	6 mg/kg every 12 hours	400 mg every 12 hours	200 mg every 12 hours
<b>Maintenance dose (after first 24 hours)</b>	4 mg/kg twice daily	200 mg twice daily	100 mg twice daily

\*This also applies to patients aged 15 years and older.

### Duration of treatment

Treatment duration should be as short as possible depending on the patient's clinical and mycological response. Long term exposure to voriconazole greater than 180 days (6 months) requires careful assessment of the benefit-risk balance (see sections 4.4 and 5.1).

### Dosage adjustment (Adults)

If patient response to treatment is inadequate, the maintenance dose may be increased to 300 mg twice daily for oral administration. For patients less than 40 kg the oral dose may be increased to 150 mg twice daily.

If patient is unable to tolerate treatment at a higher dose, reduce the oral dose by 50 mg steps to the 200 mg twice daily (or 100 mg twice daily for patients less than 40 kg) maintenance dose.

In case of use as prophylaxis, refer below.

*Children (2 to < 12 years) and young adolescents with low body weight (12 to 14 years and < 50 kg)*

Voriconazole should be dosed as children as these young adolescents may metabolize voriconazole more similarly to children than to adults.

The recommended dosing regimen is as follows:

	<b>Intravenous</b>	<b>Oral</b>
<b>Loading Dose Regimen (first 24 hours)</b>	9 mg/kg every 12 hours	Not recommended
<b>Maintenance Dose (after first 24 hours)</b>	8 mg/kg twice daily	9 mg/kg twice daily (a maximum dose of 350 mg twice daily)

Note: Based on a population pharmacokinetic analysis in 112 immunocompromised paediatric patients aged 2 to < 12 years and 26 immunocompromised adolescents aged 12 to < 17 years.

It is recommended to initiate the therapy with intravenous regimen, and oral regimen should be considered only after there is a significant clinical improvement. It should be noted that an 8 mg/kg intravenous dose will provide voriconazole exposure approximately 2-fold higher than a 9 mg/kg oral dose.

These oral dose recommendations for children are based on studies in which voriconazole was administered as the powder for oral suspension. Bioequivalence between the powder for oral suspension and tablets has not been investigated in a paediatric population. Considering the assumed limited gastro-enteric transit time in paediatric patients, the absorption of tablets may be different in paediatric compared to adult patients. It is therefore recommended to use the oral suspension formulation in children aged 2 to < 12.

*All other adolescents (12 to 14 years and  $\geq 50$  kg; 15 to 17 years regardless of body weight)*  
Voriconazole should be dosed as adults.

*Dosage adjustment (children [2 to <12 years] and young adolescents with low body weight [12 to 14 years and <50 kg])*

If patient response to treatment is inadequate, the dose may be increased by 1 mg/kg steps (or by 50 mg steps if the maximum oral dose of 350 mg was used initially). If patient is unable to tolerate treatment, reduce the dose by 1 mg/kg steps (or by 50 mg steps if the maximum oral dose of 350 mg was used initially).

Use in paediatric patients aged 2 to < 12 years with hepatic or renal insufficiency has not been studied (see sections 4.8 and 5.2).

Prophylaxis in adults and children

Prophylaxis should be initiated on the day of transplant and may be administered for up to 100 days. Prophylaxis should be as short as possible depending on the risk for developing invasive fungal infection (IFI) as defined by neutropenia or immunosuppression. It may only be continued up to 180 days after transplantation in case of continuing immunosuppression or graft versus host disease (GvHD) (see section 5.1).

### *Dosage*

The recommended dosing regimen for prophylaxis is the same as for treatment in the respective age groups. Please refer to the treatment tables above.

### *Duration of prophylaxis*

The safety and efficacy of voriconazole use for longer than 180 days has not been adequately studied in clinical trials.

Use of voriconazole in prophylaxis for greater than 180 days (6 months) requires careful assessment of the benefit-risk balance (see sections 4.4 and 5.1).

The following instructions apply to both treatment and prophylaxis

### *Dosage adjustment*

For prophylaxis use, dose adjustments are not recommended in the case of lack of efficacy or treatment-related adverse events. In the case of treatment-related adverse events, discontinuation of voriconazole and use of alternative antifungal agents must be considered (see section 4.4 and 4.8).

### *Dosage adjustments in case of co-administration*

Phenytoin may be coadministered with voriconazole if the maintenance dose of voriconazole is increased from 200 mg to 400 mg orally, twice daily (100 mg to 200 mg orally, twice daily in patients less than 40 kg), see sections 4.4 and 4.5.

The combination of voriconazole with rifabutin should, if possible be avoided. However, if the combination is strictly needed, the maintenance dose of voriconazole may be increased from 200 mg to 350 mg orally, twice daily (100 mg to 200 mg orally, twice daily in patients less than 40 kg), see sections 4.4 and 4.5.

Efavirenz may be coadministered with voriconazole if the maintenance dose of voriconazole is increased to 400 mg every 12 hours and the efavirenz dose is reduced by 50 %, i.e. to 300 mg once daily. When treatment with voriconazole is stopped, the initial dosage of efavirenz should be restored (see sections 4.4 and 4.5).

### *Elderly*

No dose adjustment is necessary for elderly patients (see section 5.2).

### *Renal impairment*

The pharmacokinetics of orally administered voriconazole are not affected by renal impairment. Therefore, no adjustment is necessary for oral dosing for patients with mild to severe renal impairment (see section 5.2).

Voriconazole is haemodialysed with a clearance of 121 ml/min. A 4-hour haemodialysis session does not remove a sufficient amount of voriconazole to warrant dose adjustment.

### *Hepatic impairment*

It is recommended that the standard loading dose regimens be used but that the maintenance dose be halved in patients with mild to moderate hepatic cirrhosis (Child-Pugh A and B) receiving voriconazole (see section 5.2).

Voriconazole has not been studied in patients with severe chronic hepatic cirrhosis (Child-Pugh C).

There is limited data on the safety of voriconazole in patients with abnormal liver function tests (aspartate transaminase [AST], alanine transaminase [ALT], alkaline phosphatase [ALP], or total bilirubin > 5 times the upper limit of normal).

Voriconazole has been associated with elevations in liver function tests and clinical signs of liver damage, such as jaundice, and must only be used in patients with severe hepatic impairment if the benefit outweighs the potential risk. Patients with severe hepatic impairment must be carefully monitored for drug toxicity (see section 4.8).

#### Paediatric population

The safety and efficacy of voriconazole in children below 2 years has not been established. Currently available data are described in sections 4.8 and 5.1 but no recommendation on a posology can be made.

#### Method of administration

<[Invented name]> film-coated tablets are to be taken at least one hour before, or one hour following, a meal.

### **4.3 Contraindications**

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

Coadministration with CYP3A4 substrates, terfenadine, astemizole, cisapride, pimozone or quinidine since increased plasma concentrations of these medicinal products can lead to QTc prolongation and rare occurrences of torsades de pointes (see section 4.5).

Coadministration with rifampicin, carbamazepine and phenobarbital since these medicinal products are likely to decrease plasma voriconazole concentrations significantly (see section 4.5).

Coadministration of standard doses of voriconazole with efavirenz doses of 400 mg once daily or higher is contraindicated, because efavirenz significantly decreases plasma voriconazole concentrations in healthy subjects at these doses. Voriconazole also significantly increases efavirenz plasma concentrations (see section 4.5, for lower doses see section 4.4).

Coadministration with high-dose ritonavir (400 mg and above twice daily) because ritonavir significantly decreases plasma voriconazole concentrations in healthy subjects at this dose (see section 4.5, for lower doses see section 4.4).

Coadministration with ergot alkaloids (ergotamine, dihydroergotamine), which are CYP3A4 substrates, since increased plasma concentrations of these medicinal products can lead to ergotism (see section 4.5).

Coadministration with sirolimus since voriconazole is likely to increase plasma concentrations of sirolimus significantly (see section 4.5).

Coadministration with St. John's Wort (see section 4.5).

#### 4.4 Special warnings and precautions for use

##### Hypersensitivity

Caution should be used in prescribing <[Invented name]> to patients with hypersensitivity to other azoles (see also section 4.8).

##### Cardiovascular

Voriconazole has been associated with QTc interval prolongation. There have been rare cases of torsades de pointes in patients taking voriconazole who had risk factors, such as history of cardiotoxic chemotherapy, cardiomyopathy, hypokalaemia and concomitant medicinal products that may have been contributory.

Voriconazole should be administered with caution to patients with potentially proarrhythmic conditions, such as:

- Congenital or acquired QTc-prolongation.
- Cardiomyopathy, in particular when heart failure is present.
- Sinus bradycardia.
- Existing symptomatic arrhythmias.
- Concomitant medicinal product that is known to prolong QTc interval. Electrolyte disturbances such as hypokalaemia, hypomagnesaemia and hypocalcaemia should be monitored and corrected, if necessary, prior to initiation and during voriconazole therapy (see section 4.2). A study has been conducted in healthy volunteers which examined the effect on QTc interval of single doses of voriconazole up to 4 times the usual daily dose. No subject experienced an interval exceeding the potentially clinically relevant threshold of 500 msec (see section 5.1).

##### Hepatic toxicity

In clinical trials, there have been cases of serious hepatic reactions during treatment with voriconazole (including clinical hepatitis, cholestasis and fulminant hepatic failure, including fatalities). Instances of hepatic reactions were noted to occur primarily in patients with serious underlying medical conditions (predominantly haematological malignancy). Transient hepatic reactions, including hepatitis and jaundice, have occurred among patients with no other identifiable risk factors. Liver dysfunction has usually been reversible on discontinuation of therapy (see section 4.8).

##### Monitoring of hepatic function

Patients receiving voriconazole must be carefully monitored for hepatic toxicity. Clinical management should include laboratory evaluation of hepatic function (specifically AST and ALT) at the initiation of treatment with voriconazole and at least weekly for the first month of treatment. Treatment duration should be as short as possible; however, if based on the benefit-risk assessment the treatment is continued (see section 4.2), monitoring frequency can be reduced to monthly if there are no changes in the liver function tests.

If the liver function tests become markedly elevated, <[Invented name]> should be discontinued, unless the medical judgment of the risk-benefit of the treatment for the patient justifies continued use.

Monitoring of hepatic function should be carried out in both children and adults.

### Serious dermatological adverse reactions

#### Phototoxicity

In addition voriconazole has been associated with phototoxicity, including reactions such as ephelides, lentigo, actinic keratosis and pseudoporphyria. It is recommended that all patients, including children, avoid exposure to direct sunlight during <[Invented name]> treatment and use measures such as protective clothing and sunscreen with high sun protection factor (SPF).

#### Squamous cell carcinoma of the skin (SCC)

Squamous cell carcinoma of the skin has been reported in patients, some of whom have reported prior phototoxic reactions. If phototoxic reactions occur, multidisciplinary advice should be sought, <[Invented name]> discontinuation and use of alternative antifungal agents should be considered and the patient should be referred to a dermatologist. If voriconazole is continued, however, dermatologic evaluation should be performed on a systematic and regular basis, to allow early detection and management of premalignant lesions. <[Invented name]> should be discontinued if premalignant skin lesions or squamous cell carcinoma are identified (see below the section under Long-term treatment).

#### Exfoliative cutaneous reactions

Severe cutaneous adverse reactions (SCARs) including Stevens-Johnson syndrome (SJS), toxic epidermal necrolysis (TEN), and drug reaction with eosinophilia and systematic symptoms (DRESS), which can be life-threatening or fatal, have been reported with the use voriconazole. If a patient develops a rash he should be monitored closely and <[Invented name]> discontinued if lesions progress.

### Long-term treatment

Long term exposure (treatment or prophylaxis) greater than 180 days (6 months) requires careful assessment of the benefit-risk balance and physicians should therefore consider the need to limit the exposure to <[Invented name]> (see sections 4.2 and 5.1).

Squamous cell carcinoma of the skin (SCC) has been reported in relation with long-term voriconazole treatment.

Non-infectious periostitis with elevated fluoride and alkaline phosphatase levels has been reported in transplant patients. If a patient develops skeletal pain and radiologic findings compatible with periostitis <[Invented name]> discontinuation should be considered after multidisciplinary advice.

### Visual adverse reactions

There have been reports of prolonged visual adverse reactions, including blurred vision, optic neuritis and papilloedema (see section 4.8).

### Renal adverse reactions

Acute renal failure has been observed in severely ill patients undergoing treatment with voriconazole. Patients being treated with voriconazole are likely to be treated concomitantly with nephrotoxic medicinal products and have concurrent conditions that may result in decreased renal function (see section 4.8).

#### Monitoring of renal function

Patients should be monitored for the development of abnormal renal function. This should include laboratory evaluation, particularly serum creatinine.

#### Monitoring of pancreatic function

Patients, especially children, with risk factors for acute pancreatitis (e.g., recent chemotherapy, haematopoietic stem cell transplantation [HSCT]), should be monitored closely during <[Invented name]> treatment. Monitoring of serum amylase or lipase may be considered in this clinical situation.

#### Paediatric population

Safety and effectiveness in paediatric subjects below the age of two years has not been established (see sections 4.8 and 5.1). Voriconazole is indicated for paediatric patients aged two years or older. A higher frequency of liver enzyme elevations was observed in the paediatric population (see section 4.8). Hepatic function should be monitored in both children and adults. Oral bioavailability may be limited in paediatric patients aged 2 to < 12 years with malabsorption and very low body weight for age. In that case, intravenous voriconazole administration is recommended.

#### *Serious dermatological adverse reactions (including SCC)*

The frequency of phototoxicity reactions is higher in the paediatric population. As an evolution towards SCC has been reported, stringent measures for the photoprotection are warranted in this population of patients. In children experiencing photoaging injuries such as lentigines or ephelides, sun avoidance and dermatologic follow-up are recommended even after treatment discontinuation.

#### Prophylaxis

In case of treatment-related adverse events (hepatotoxicity, severe skin reactions including phototoxicity and SCC, severe or prolonged visual disorders and periostitis), discontinuation of voriconazole and use of alternative antifungal agents must be considered.

#### Phenytoin (CYP2C9 substrate and potent CYP450 inducer)

Careful monitoring of phenytoin levels is recommended when phenytoin is coadministered with voriconazole. Concomitant use of voriconazole and phenytoin should be avoided unless the benefit outweighs the risk (see section 4.5).

#### Efavirenz (CYP450 inducer; CYP3A4 inhibitor and substrate)

When voriconazole is coadministered with efavirenz the dose of voriconazole should be increased to 400 mg every 12 hours and the dose of efavirenz should be decreased to 300 mg every 24 hours (see sections 4.2, 4.3 and 4.5).

#### Rifabutin (potent CYP450 inducer)



Careful monitoring of full blood counts and adverse reactions to rifabutin (e.g., uveitis) is recommended when rifabutin is coadministered with voriconazole. Concomitant use of voriconazole and rifabutin should be avoided unless the benefit outweighs the risk (see section 4.5).

#### Ritonavir (potent CYP450 inducer; CYP3A4 inhibitor and substrate)

Coadministration of voriconazole and low dose ritonavir (100 mg twice daily) should be avoided unless an assessment of the benefit/risk to the patient justifies the use of voriconazole (see sections 4.3 and 4.5).

#### Everolimus (CYP3A4 substrate, P-gp substrate)

Coadministration of voriconazole with everolimus is not recommended because voriconazole is expected to significantly increase everolimus concentrations. Currently there are insufficient data to allow dosing recommendations in this situation (see section 4.5).

#### Methadone (CYP3A4 substrate)

Frequent monitoring for adverse reactions and toxicity related to methadone, including QTc prolongation, is recommended when coadministered with voriconazole since methadone levels increased following coadministration of voriconazole. Dose reduction of methadone may be needed (see section 4.5).

#### Short-acting opiates (CYP3A4 substrate)

Reduction in the dose of alfentanil, fentanyl and other short-acting opiates similar in structure to alfentanil and metabolised by CYP3A4 (e.g., sufentanil) should be considered when coadministered with voriconazole (see section 4.5). As the half-life of alfentanil is prolonged in a 4-fold manner when alfentanil is coadministered with voriconazole, and in an independent published study concomitant use of voriconazole with fentanyl resulted in an increase in the mean AUC<sub>0-∞</sub> of fentanyl, frequent monitoring for opiate-associated adverse reactions (including a longer respiratory monitoring period) may be necessary.

#### Long-acting opiates (CYP3A4 substrate)

Reduction in the dose of oxycodone and other long-acting opiates metabolized by CYP3A4 (e.g., hydrocodone) should be considered when coadministered with voriconazole. Frequent monitoring for opiate-associated adverse reactions may be necessary (see section 4.5).

#### Fluconazole (CYP2C9, CYP2C19 and CYP3A4 inhibitor)

Coadministration of oral voriconazole and oral fluconazole resulted in a significant increase in C<sub>max</sub> and AUC<sub>τ</sub> of voriconazole in healthy subjects. The reduced dose and/or frequency of voriconazole and fluconazole that would eliminate this effect have not been established. Monitoring for voriconazole-associated adverse reactions is recommended if voriconazole is used sequentially after fluconazole (see section 4.5).

<[Invented name]> film-coated tablets contain lactose. Patients with rare hereditary problems of galactose intolerance, total lactase deficiency or glucose-galactose malabsorption should not take this medicine.

### **4.5 Interaction with other medicinal products and other forms of interaction**

Voriconazole is metabolised by, and inhibits the activity of, cytochrome P450 isoenzymes, CYP2C19, CYP2C9, and CYP3A4. Inhibitors or inducers of these isoenzymes may increase or decrease voriconazole plasma concentrations, respectively, and there is potential for voriconazole to increase the plasma concentrations of substances metabolised by these CYP450 isoenzymes.

Unless otherwise specified, drug interaction studies have been performed in healthy adult male subjects using multiple dosing to steady state with oral voriconazole at 200 mg twice daily (BID). These results are relevant to other populations and routes of administration.

Voriconazole should be administered with caution in patients with concomitant medication that is known to prolong QTc interval. When there is also a potential for voriconazole to increase the plasma concentrations of substances metabolised by CYP3A4 isoenzymes (certain antihistamines, quinidine, cisapride, pimozone) coadministration is contraindicated (see below and section 4.3).

#### Interaction table

Interactions between voriconazole and other medicinal products are listed in the table below (once daily as “QD”, twice daily as “BID”, three times daily as “TID” and not determined as “ND”). The direction of the arrow for each pharmacokinetic parameter is based on the 90 % confidence interval of the geometric mean ratio being within ( $\leftrightarrow$ ), below ( $\downarrow$ ) or above ( $\uparrow$ ) the 80-125 % range. The asterisk (\*) indicates a two-way interaction.  $AUC_{\tau}$ ,  $AUC_t$  and  $AUC_{0-\infty}$  represent area under the curve over a dosing interval, from time zero to the time with detectable measurement and from time zero to infinity, respectively.

The interactions in the table are presented in the following order: contraindications, those requiring dose adjustment and careful clinical and/or biological monitoring, and finally those that have no significant pharmacokinetic interaction but may be of clinical interest in this therapeutic field.

<b>Medicinal product</b> <i>[Mechanism of Interaction]</i>	<b>Interaction</b> <b>Geometric mean changes (%)</b>	<b>Recommendations</b> <b>concerning</b> <b>coadministration</b>
Astemizole, cisapride, pimozone, quinidine and terfenadine <i>[CYP3A4 substrates]</i>	Although not studied, increased plasma concentrations of these medicinal products can lead to QTc prolongation and rare occurrences of torsades de pointes.	<b>Contraindicated</b> (see section 4.3)
Carbamazepine and long-acting barbiturates (e.g., phenobarbital, mephobarbital) <i>[potent CYP450 inducers]</i>	Although not studied, carbamazepine and long-acting barbiturates are likely to significantly decrease plasma voriconazole concentrations.	<b>Contraindicated</b> (see section 4.3)
Efavirenz (a non-nucleoside reverse transcriptase inhibitor) <i>[CYP450 inducer; CYP3A4 inhibitor and substrate]</i>  Efavirenz 400 mg QD,	   Efavirenz $C_{max}$ $\uparrow$ 38%	   Use of standard doses of voriconazole with efavirenz doses of 400 mg QD or higher

<p>coadministered with voriconazole 200 mg BID*</p> <p>Efavirenz 300 mg QD, co-administered with voriconazole 400 mg BID*</p>	<p>Efavirenz <math>AUC_{\tau}</math> ↑ 44%                  Voriconazole <math>C_{max}</math> ↓ 61%                  Voriconazole <math>AUC_{\tau}</math> ↓ 77%</p> <p>Compared to efavirenz 600 mg QD,                  Efavirenz <math>C_{max}</math> ↔                  Efavirenz <math>AUC_{\tau}</math> ↑ 17%</p> <p>Compared to voriconazole 200 mg BID,                  Voriconazole <math>C_{max}</math> ↑ 23%                  Voriconazole <math>AUC_{\tau}</math> ↓ 7%</p>	<p>is <b>contraindicated</b> (see section 4.3).</p> <p>Voriconazole may be co-administered with efavirenz if the voriconazole maintenance dose is increased to 400 mg BID and the efavirenz dose is decreased to 300 mg QD. When voriconazole treatment is stopped, the initial dose of efavirenz should be restored (see sections 4.2 and 4.4).</p>
<p>Ergot alkaloids (e.g., ergotamine and dihydroergotamine)                  [CYP3A4 substrates]</p>	<p>Although not studied, voriconazole is likely to increase the plasma concentrations of ergot alkaloids and lead to ergotism.</p>	<p><b>Contraindicated</b> (see section 4.3)</p>
<p>Rifabutin                  [potent CYP450 inducer]</p> <p>300 mg QD</p> <p>300 mg QD (co-administered with voriconazole 350 mg BID)*</p> <p>300 mg QD (co-administered with voriconazole 400 mg BID)*</p>	<p>Voriconazole <math>C_{max}</math> ↓ 69%                  Voriconazole <math>AUC_{\tau}</math> ↓ 78%</p> <p>Compared to voriconazole 200 mg BID,                  Voriconazole <math>C_{max}</math> ↓ 4%                  Voriconazole <math>AUC_{\tau}</math> ↓ 32%</p> <p>Rifabutin <math>C_{max}</math> ↑ 195%                  Rifabutin <math>AUC_{\tau}</math> ↑ 331%</p> <p>Compared to voriconazole 200 mg BID,                  Voriconazole <math>C_{max}</math> ↑ 104%                  Voriconazole <math>AUC_{\tau}</math> ↑ 87%</p>	<p>Concomitant use of voriconazole and rifabutin should be avoided unless the benefit outweighs the risk. The maintenance dose of voriconazole may be increased to 5 mg/kg intravenously BID or from 200 mg to 350 mg orally BID (100 mg to 200 mg orally BID in patients less than 40 kg) (see section 4.2). Careful monitoring of full blood counts and adverse reactions to rifabutin (e.g., uveitis) is recommended when rifabutin is coadministered with voriconazole.</p>
<p>Rifampicin (600 mg QD)                  [potent CYP450 inducer]</p>	<p>Voriconazole <math>C_{max}</math> ↓ 93%                  Voriconazole <math>AUC_{\tau}</math> ↓ 96%</p>	<p><b>Contraindicated</b> (see section 4.3)</p>
<p>Ritonavir (protease inhibitor)                  [potent CYP450 inducer; CYP3A4 inhibitor and substrate]</p> <p>High dose (400 mg BID)</p>	<p>Ritonavir <math>C_{max}</math> and <math>AUC_{\tau}</math> ↔                  Voriconazole <math>C_{max}</math> ↓ 66%</p>	<p>Co-administration of voriconazole and high doses of ritonavir (400 mg and above</p>

<p>Low dose (100 mg BID)*</p>	<p>Voriconazole AUC<sub>τ</sub> ↓ 82%</p> <p>Ritonavir C<sub>max</sub> ↓ 25%</p> <p>Ritonavir AUC<sub>τ</sub> ↓ 13%</p> <p>Voriconazole C<sub>max</sub> ↓ 24%</p> <p>Voriconazole AUC<sub>τ</sub> ↓ 39%</p>	<p>BID) is <b>contraindicated</b> (see section 4.3).</p> <p>Coadministration of voriconazole and low dose ritonavir (100 mg BID) should be avoided, unless an assessment of the benefit/risk to the patient justifies the use of voriconazole.</p>
<p>St. John's Wort  <i>[CYP450 inducer; P-gp inducer]</i></p> <p>300 mg TID (co-administered with voriconazole 400 mg single dose)</p>	<p>In an independent published study,                  Voriconazole AUC<sub>0-∞</sub> ↓ 59%</p>	<p><b>Contraindicated</b> (see section 4.3)</p>
<p>Everolimus  <i>[CYP3A4 substrate, P-gp substrate]</i></p>	<p>Although not studied, voriconazole is likely to significantly increase the plasma concentrations of everolimus.</p>	<p>Co-administration of voriconazole with everolimus is not recommended because voriconazole is expected to significantly increase everolimus concentrations (see section 4.4).</p>
<p>Fluconazole (200 mg QD)  <i>[CYP2C9, CYP2C19 and CYP3A4 inhibitor]</i></p>	<p>Voriconazole C<sub>max</sub> ↑ 57%</p> <p>Voriconazole AUC<sub>τ</sub> ↑ 79%</p> <p>Fluconazole C<sub>max</sub> ND</p> <p>Fluconazole AUC<sub>τ</sub> ND</p>	<p>The reduced dose and/or frequency of voriconazole and fluconazole that would eliminate this effect have not been established. Monitoring for voriconazole-associated adverse reactions is recommended if voriconazole is used sequentially after fluconazole.</p>
<p>Phenytoin  <i>[CYP2C9 substrate and potent CYP450 inducer]</i></p> <p>300 mg QD</p> <p>300 mg QD (co-administered with voriconazole 400 mg BID)*</p>	<p>Voriconazole C<sub>max</sub> ↓ 49%</p> <p>Voriconazole AUC<sub>τ</sub> ↓ 69%</p> <p>Phenytoin C<sub>max</sub> ↑ 67%</p> <p>Phenytoin AUC<sub>τ</sub> ↑ 81%</p> <p>Compared to voriconazole 200 mg BID,                  Voriconazole C<sub>max</sub> ↑ 34%</p> <p>Voriconazole AUC<sub>τ</sub> ↑ 39%</p>	<p>Concomitant use of voriconazole and phenytoin should be avoided unless the benefit outweighs the risk. Careful monitoring of phenytoin plasma levels is recommended.</p> <p>Phenytoin may be co-administered with voriconazole if the maintenance dose of voriconazole is increased to 5 mg/kg IV BID or from 200 mg to 400 mg oral BID, (100 mg to 200 mg oral BID in patients less than 40 kg) (see section 4.2).</p>
<p>Anticoagulants</p>		<p>Close monitoring of prothrombin time or other</p>

<p>Warfarin (30 mg single dose, co-administered with 300 mg BID voriconazole)  <i>[CYP2C9 substrate]</i></p> <p>Other oral coumarins (e.g., phenprocoumon, acenocoumarol)  <i>[CYP2C9 and CYP3A4 substrates]</i></p>	<p>Maximum increase in prothrombin time was approximately 2-fold.</p> <p>Although not studied, voriconazole may increase the plasma concentrations of coumarins that may cause an increase in prothrombin time.</p>	<p>suitable anticoagulation tests is recommended, and the dose of anticoagulants should be adjusted accordingly.</p>
<p>Benzodiazepines (e.g., midazolam, triazolam, alprazolam)  <i>[CYP3A4 substrates]</i></p>	<p>Although not studied clinically, voriconazole is likely to increase the plasma concentrations of benzodiazepines that are metabolised by CYP3A4 and lead to a prolonged sedative effect.</p>	<p>Dose reduction of benzodiazepines should be considered.</p>
<p>Immunosuppressants  <i>[CYP3A4 substrates]</i></p> <p>Sirolimus (2 mg single dose)</p> <p>Ciclosporin (in stable renal transplant recipients receiving chronic ciclosporin therapy)</p> <p>Tacrolimus (0.1 mg/kg single dose)</p>	<p>In an independent published study,              Sirolimus <math>C_{max}</math> ↑ 6.6-fold              Sirolimus <math>AUC_{0-\infty}</math> ↑ 11-fold</p> <p>Ciclosporin <math>C_{max}</math> ↑ 13%              Ciclosporin <math>AUC_{\tau}</math> ↑ 70%</p> <p>Tacrolimus <math>C_{max}</math> ↑ 117%              Tacrolimus <math>AUC_{\tau}</math> ↑ 221%</p>	<p>Coadministration of voriconazole and sirolimus is <b>contraindicated</b> (see section 4.3).</p> <p>When initiating voriconazole in patients already on ciclosporin it is recommended that the ciclosporin dose be halved and ciclosporin level carefully monitored. Increased ciclosporin levels have been associated with nephrotoxicity. <u>When voriconazole is discontinued, ciclosporin levels must be carefully monitored and the dose increased as necessary.</u></p> <p>When initiating voriconazole in patients already on tacrolimus, it is recommended that the tacrolimus dose be reduced to a third of the original dose and tacrolimus level carefully monitored. Increased tacrolimus levels have been associated with nephrotoxicity. <u>When voriconazole is discontinued, tacrolimus levels</u></p>

		<u>must be carefully monitored and the dose increased as necessary.</u>
Long Acting Opiates [CYP3A4 substrates]  Oxycodone (10 mg single dose)	In an independent published study, Oxycodone $C_{max}$ ↑ 1.7-fold Oxycodone $AUC_{0-\infty}$ ↑ 3.6-fold	Dose reduction in oxycodone and other long-acting opiates metabolized by CYP3A4 (e.g., hydrocodone) should be considered. Frequent monitoring for opiate-associated adverse reactions may be necessary.
Methadone (32-100 mg QD) [CYP3A4 substrate]	R-methadone (active) $C_{max}$ ↑ 31% R-methadone (active) $AUC_{\tau}$ ↑ 47% S-methadone $C_{max}$ ↑ 65% S-methadone $AUC_{\tau}$ ↑ 103%	Frequent monitoring for adverse reactions and toxicity related to methadone, including QTc prolongation, is recommended. Dose reduction of methadone may be needed.
Non-Steroidal Anti-Inflammatory Drugs (NSAIDs) [CYP2C9 substrates]  Ibuprofen (400 mg single dose)  Diclofenac (50 mg single dose)	S-Ibuprofen $C_{max}$ ↑ 20% S-Ibuprofen $AUC_{0-\infty}$ ↑ 100%  Diclofenac $C_{max}$ ↑ 114% Diclofenac $AUC_{0-\infty}$ ↑ 78%	Frequent monitoring for adverse reactions and toxicity related to NSAIDs is recommended. Dose reduction of NSAIDs may be needed.
Omeprazole (40 mg QD)* [CYP2C19 inhibitor; CYP2C19 and CYP3A4 substrate]	Omeprazole $C_{max}$ ↑ 116% Omeprazole $AUC_{\tau}$ ↑ 280% Voriconazole $C_{max}$ ↑ 15% Voriconazole $AUC_{\tau}$ ↑ 41%  Other proton pump inhibitors that are CYP2C19 substrates may also be inhibited by voriconazole and may result in increased plasma concentrations of these medicinal products.	No dose adjustment of voriconazole is recommended.  When initiating voriconazole in patients already receiving omeprazole doses of 40 mg or above, it is recommended that the omeprazole dose be halved.
Oral Contraceptives* [CYP3A4 substrate; CYP2C19 inhibitor]  Norethisterone/ethinylestradiol (1 mg/0.035 mg QD)	Ethinylestradiol $C_{max}$ ↑ 36% Ethinylestradiol $AUC_{\tau}$ ↑ 61% Norethisterone $C_{max}$ ↑ 15% Norethisterone $AUC_{\tau}$ ↑ 53% Voriconazole $C_{max}$ ↑ 14% Voriconazole $AUC_{\tau}$ ↑ 46%	Monitoring for adverse reactions related to oral contraceptives, in addition to those for voriconazole, is recommended.
Short Acting Opiates [CYP3A4 substrates]  Alfentanil (20 µg/kg single dose, with concomitant naloxone)	In an independent published study, Alfentanil $AUC_{0-\infty}$ ↑ 6-fold	Dose reduction of alfentanil, fentanyl and other short acting opiates similar in structure to alfentanil and metabolised by CYP3A4 (e.g., sufentanil) should be considered. Extended and frequent monitoring for respiratory depression and other

Fentanyl (5 µg/kg single dose)	In an independent published study, Fentanyl AUC <sub>0-∞</sub> ↑ 1.34-fold	opiate- associated adverse reactions is recommended.
Statins (e.g., lovastatin) [CYP3A4 substrates]	Although not studied clinically, voriconazole is likely to increase the plasma concentrations of statins that are metabolised by CYP3A4 and could lead to rhabdomyolysis.	Dose reduction of statins should be considered.
Sulphonylureas (e.g., tolbutamide, glipizide, glyburide) [CYP2C9 substrates]	Although not studied, voriconazole is likely to increase the plasma concentrations of sulphonylureas and cause hypoglycaemia.	Careful monitoring of blood glucose is recommended. Dose reduction of sulphonylureas should be considered.
Vinca Alkaloids (e.g., vincristine and vinblastine) [CYP3A4 substrates]	Although not studied, voriconazole is likely to increase the plasma concentrations of vinca alkaloids and lead to neurotoxicity.	Dose reduction of vinca alkaloids should be considered.
Other HIV Protease Inhibitors (e.g., saquinavir, amprenavir and nelfinavir)* [CYP3A4 substrates and inhibitors]	Not studied clinically. <i>In vitro</i> studies show that voriconazole may inhibit the metabolism of HIV protease inhibitors and the metabolism of voriconazole may also be inhibited by HIV protease inhibitors.	Careful monitoring for any occurrence of drug toxicity and/or lack of efficacy, and dose adjustment may be needed.
Other Non-Nucleoside Reverse Transcriptase Inhibitors (NNRTIs) (e.g., delavirdine, nevirapine)* [CYP3A4 substrates, inhibitors or CYP450 inducers]	Not studied clinically. <i>In vitro</i> studies show that the metabolism of voriconazole may be inhibited by NNRTIs and voriconazole may inhibit the metabolism of NNRTIs. The findings of the effect of efavirenz on voriconazole suggest that the metabolism of voriconazole may be induced by a NNRTI.	Careful monitoring for any occurrence of drug toxicity and/or lack of efficacy, and dose adjustment may be needed.
Cimetidine (400 mg BID) [non-specific CYP450 inhibitor and increases gastric pH]	Voriconazole C <sub>max</sub> ↑ 18% Voriconazole AUC <sub>τ</sub> ↑ 23%	No dose adjustment
Digoxin (0.25 mg QD) [P-gp substrate]	Digoxin C <sub>max</sub> ↔ Digoxin AUC <sub>τ</sub> ↔	No dose adjustment
Indinavir (800 mg TID) [CYP3A4 inhibitor and substrate]	Indinavir C <sub>max</sub> ↔ Indinavir AUC <sub>τ</sub> ↔ Voriconazole C <sub>max</sub> ↔ Voriconazole AUC <sub>τ</sub> ↔	No dose adjustment
Macrolide antibiotics		No dose adjustment

Erythromycin (1 g BID) [CYP3A4 inhibitor]	Voriconazole C <sub>max</sub> and AUC <sub>τ</sub> ↔	
Azithromycin (500 mg QD)	Voriconazole C <sub>max</sub> and AUC <sub>τ</sub> ↔  The effect of voriconazole on either erythromycin or azithromycin is unknown.	
Mycophenolic acid (1 g single dose) [UDP-glucuronyl transferase substrate]	Mycophenolic acid C <sub>max</sub> ↔ Mycophenolic acid AUC <sub>t</sub> ↔	No dose adjustment
Prednisolone (60 mg single dose) [CYP3A4 substrate]	Prednisolone C <sub>max</sub> ↑ 11% Prednisolone AUC <sub>0-∞</sub> ↑ 34%	No dose adjustment
Ranitidine (150 mg BID) [increases gastric pH]	Voriconazole C <sub>max</sub> and AUC <sub>τ</sub> ↔	No dose adjustment

#### 4.6 Fertility, pregnancy and lactation

##### Pregnancy

There are no adequate data on the use of voriconazole in pregnant women available.

Studies in animals have shown reproductive toxicity (see section 5.3). The potential risk for humans is unknown.

<[Invented name]> must not be used during pregnancy unless the benefit to the mother clearly outweighs the potential risk to the foetus.

##### Women of child-bearing potential

Women of child-bearing potential must always use effective contraception during treatment.

##### Breast-feeding

The excretion of voriconazole into breast milk has not been investigated. Breast-feeding must be stopped on initiation of treatment with <[Invented name]>.

##### Fertility

In an animal study, no impairment of fertility was demonstrated in male and female rats (see section 5.3).

#### 4.7 Effects on ability to drive and use machines

<[Invented name]> has moderate influence on the ability to drive and use machines. It may cause transient and reversible changes to vision, including blurring, altered/enhanced visual perception and/or photophobia. Patients must avoid potentially hazardous tasks, such as driving or operating machinery while experiencing these symptoms.

#### 4.8 Undesirable effects



Summary of safety profile

The safety profile of voriconazole in adults is based on an integrated safety database of more than 2,000 subjects (including 1,603 adult patients in therapeutic trials) and an additional 270 adults in prophylaxis trials. This represents a heterogeneous population, containing patients with haematological malignancy, HIV-infected patients with oesophageal candidiasis and refractory fungal infections, non-neutropenic patients with candidaemia or aspergillosis and healthy volunteers.

The most commonly reported adverse reactions were visual impairment, pyrexia, rash, vomiting, nausea, diarrhoea, headache, peripheral oedema, liver function test abnormal, respiratory distress and abdominal pain.

The severity of the adverse reactions was generally mild to moderate. No clinically significant differences were seen when the safety data were analysed by age, race, or gender.

Tabulated list of adverse reactions

In the table below, since the majority of the studies were of an open nature, all causality adverse reactions and their frequency categories in 1,873 adults from pooled therapeutic (1,603) and prophylaxis (270) studies, by system organ class, are listed.

Frequency categories are expressed as: Very common ( $\geq 1/10$ ); Common ( $\geq 1/100$  to  $< 1/10$ ); Uncommon ( $\geq 1/1,000$  to  $< 1/100$ ); Rare ( $\geq 1/10,000$  to  $< 1/1,000$ ); Very rare ( $< 1/10,000$ ); Not known (cannot be estimated from the available data).

Within each frequency grouping, undesirable effects are presented in order of decreasing seriousness.

Undesirable effects reported in subjects receiving voriconazole:

<b>System Organ Class</b>	<b>Very common <math>\geq 1/10</math></b>	<b>Common <math>\geq 1/100</math> to <math>&lt; 1/10</math></b>	<b>Uncommon <math>\geq 1/1,000</math> to <math>&lt; 1/100</math></b>	<b>Rare <math>\geq 1/10,000</math> to <math>&lt; 1/1,000</math></b>	<b>Frequency not known (cannot be estimated from available data)</b>
<b>Infections and infestations</b>		sinusitis	pseudomembranous colitis		
<b>Neoplasms benign, malignant and unspecified (including cysts and polyps)</b>					squamous cell carcinoma*

System Organ Class	Very common ≥ 1/10	Common ≥ 1/100 to < 1/10	Uncommon ≥ 1/1,000 to < 1/100	Rare ≥ 1/10,000 to < 1/1,000	Frequency not known (cannot be estimated from available data)
<b>Blood and lymphatic system disorders</b>		agranulocytosis <sup>1</sup> , pancytopenia, thrombocytopenia <sup>2</sup> , leukopenia, anaemia	bone marrow failure, lymphadenopathy, eosinophilia	disseminated intravascular coagulation	
<b>Immune system disorders</b>			hypersensitivity	anaphylactoid reaction	
<b>Endocrine disorders</b>			adrenal insufficiency, hypothyroidism	hyperthyroidism	
<b>Metabolism and nutrition disorders</b>	oedema peripheral	hypoglycaemia, hypokalaemia, hyponatraemia			
<b>Psychiatric disorders</b>		depression, hallucination, anxiety, insomnia, agitation, confusional state			
<b>Nervous system disorders</b>	headache	convulsion, syncope, tremor, hypertonia <sup>3</sup> , paraesthesia, somnia, dizziness	brain oedema, encephalopathy <sup>4</sup> , extrapyramidal disorder <sup>5</sup> , neuropathy peripheral, ataxia, hypoesthesia, dysgeusia,	hepatic encephalopathy, Guillain-Barre syndrome, nystagmus	
<b>Eye disorders</b>	visual impairment <sup>6</sup>	retinal haemorrhage	oculogyric crisis, optic nerve disorder <sup>7</sup> , papilloedema <sup>8</sup> , scleritis, blepharitis, diplopia	optic atrophy, corneal opacity	

System Organ Class	Very common ≥ 1/10	Common ≥ 1/100 to < 1/10	Uncommon ≥ 1/1,000 to < 1/100	Rare ≥ 1/10,000 to < 1/1,000	Frequency not known (cannot be estimated from available data)
<b>Ear and labyrinth disorders</b>			hypoacusis, vertigo, tinnitus		
<b>Cardiac disorders</b>		arrhythmia supraventricular, tachycardia, bradycardia	Ventricular fibrillation, ventricular extrasystoles, supraventricular tachycardia, ventricular tachycardia, electrocardiogram QT prolonged	torsades de pointes, atrioventricular block complete, bundle branch block, nodal rhythm	
<b>Vascular disorders</b>		hypotension, phlebitis	thrombophlebitis, lymphangitis		
<b>Respiratory, thoracic and mediastinal disorders</b>	respiratory distress <sup>9</sup>	acute respiratory distress syndrome, pulmonary oedema			
<b>Gastrointestinal disorders</b>	diarrhoea, vomiting, abdominal pain, nausea	cheilitis, dyspepsia, constipation, gingivitis	peritonitis, pancreatitis, swollen tongue, duodenitis, gastroenteritis, glossitis		
<b>Hepatobiliary disorders</b>	liver function test abnormal	jaundice, jaundice cholestatic, hepatitis <sup>10</sup>	hepatic failure, hepatomegaly, cholecystitis, cholelithiasis		
<b>Skin and subcutaneous tissue disorders</b>	rash	dermatitis exfoliative, alopecia, rash maculopapular, pruritus, erythema	Stevens-Johnson syndrome (SJS), urticaria, dermatitis allergic,	toxic epidermal necrolysis (TEN), angioedema, actinic keratosis*,	cutaneous lupus erythematosus*, ephelides*, lentigo*

System Organ Class	Very common ≥ 1/10	Common ≥ 1/100 to < 1/10	Uncommon ≥ 1/1,000 to < 1/100	Rare ≥ 1/10,000 to < 1/1,000	Frequency not known (cannot be estimated from available data)
			phototoxicity, rash macular, rash papular, purpura, eczema	pseudoporphyria, erythema multiforme, psoriasis, drug eruption, drug reaction with eosinophilia and systematic symptoms (DRESS)	
<b>Musculoskeletal and connective tissue disorders</b>		back pain	arthritis		periostitis*
<b>Renal and urinary disorders</b>		renal failure acute, haematuria	renal tubular necrosis, proteinuria, nephritis		
<b>General disorders and administration site conditions</b>	pyrexia	chest pain, face oedema <sup>11</sup> , asthenia, chills	infusion site reaction, influenza like illness		
<b>Investigations</b>		blood creatinine increased	blood urea increased, blood cholesterol increased		

\*ADR identified post-marketing

<sup>1</sup> Includes febrile neutropenia and neutropenia.

<sup>2</sup> Includes immune thrombocytopenic purpura.

<sup>3</sup> Includes nuchal rigidity and tetany.

<sup>4</sup> Includes hypoxic-ischaemic encephalopathy and metabolic encephalopathy.

<sup>5</sup> Includes akathisia and parkinsonism.

<sup>6</sup> See “Visual impairments” paragraph in section 4.8.

<sup>7</sup> Prolonged optic neuritis has been reported post-marketing. See section 4.4.

<sup>8</sup> See section 4.4.

<sup>9</sup> Includes dyspnoea and dyspnoea exertional.

<sup>10</sup> Includes drug-induced liver injury, hepatitis toxic, hepatocellular injury and hepatotoxicity.

<sup>11</sup> Includes periorbital oedema, lip oedema, and oedema mouth.

### Description of selected adverse reactions

#### *Visual impairments*

In clinical trials, visual impairments (including blurred vision, photophobia, chloropsia, chromatopsia, colour blindness, cyanopsia, eye disorder, halo vision, night blindness, oscillopsia, photopsia, scintillating scotoma, visual acuity reduced, visual brightness, visual field defect, vitreous floaters, and xanthopsia) with voriconazole were very common. These visual impairments were transient and fully reversible, with the majority spontaneously resolving within 60 minutes and no clinically significant long-term visual effects were observed. There was evidence of attenuation with repeated doses of voriconazole. The visual impairments were generally mild, rarely resulted in discontinuation and were not associated with long-term sequelae. Visual impairments may be associated with higher plasma concentrations and/or doses.

The mechanism of action is unknown, although the site of action is most likely to be within the retina. In a study in healthy volunteers investigating the impact of voriconazole on retinal function, voriconazole caused a decrease in the electroretinogram (ERG) waveform amplitude. The ERG measures electrical currents in the retina. The ERG changes did not progress over 29 days of treatment and were fully reversible on withdrawal of voriconazole.

There have been post-marketing reports of prolonged visual adverse events (see section 4.4).

#### *Dermatological reactions*

Dermatological reactions were very common in patients treated with voriconazole in clinical trials, but these patients had serious underlying diseases and were receiving multiple concomitant medicinal products. The majority of rashes were of mild to moderate severity. Patients have developed severe cutaneous adverse reactions (SCARs), including Stevens-Johnson syndrome (SJS) (uncommon), toxic epidermal necrolysis (TEN) (rare), drug reaction with eosinophilia and systematic symptoms (DRESS) (rare) and erythema multiforme (rare) during treatment with voriconazole (see section 4.4).

If a patient develops a rash they should be monitored closely and <[Invented name]> discontinued if lesions progress.

Photosensitivity reactions such as ephelides, lentigo and actinic keratosis have been reported, especially during long-term therapy (see section 4.4).

There have been reports of squamous cell carcinoma of the skin in patients treated with voriconazole for long periods of time; the mechanism has not been established (see section 4.4).

#### *Liver function tests*

The overall incidence of transaminase increases >3 xULN (not necessarily comprising an adverse event) in the voriconazole clinical programme was 18 % (319/1,768) in adults and 25.8% (73/283) in paediatric subjects who received voriconazole for pooled therapeutic and

prophylaxis use. Liver function test abnormalities may be associated with higher plasma concentrations and/or doses.

The majority of abnormal liver function tests either resolved during treatment without dose adjustment or following dose adjustment, including discontinuation of therapy.

Voriconazole has been associated with cases of serious hepatic toxicity in patients with other serious underlying conditions. This includes cases of jaundice, hepatitis and hepatic failure leading to death (see section 4.4).

#### *Prophylaxis*

In an open-label, comparative, multicenter study comparing voriconazole and itraconazole as primary prophylaxis in adult and adolescent allogeneic HSCT recipients without prior proven or probable IFI, permanent discontinuation of voriconazole due to AEs was reported in 39.3% of subjects versus 39.6% of subjects in the itraconazole arm. Treatment-emergent hepatic AEs resulted in permanent discontinuation of study medication for 50 subjects (21.4%) treated with voriconazole and for 18 subjects (7.1%) treated with itraconazole.

#### *Paediatric population*

The safety of voriconazole was investigated in 288 paediatric patients aged 2 to < 12 years (169) and 12 to <18 years (119) who received voriconazole for prophylaxis (183) and therapeutic use (105) in clinical trials. The safety of voriconazole was also investigated in 158 additional paediatric patients aged 2 to <12 years in compassionate use programs. Overall, the safety profile of voriconazole in paediatric population was similar to that in adults. However, a trend towards a higher frequency of liver enzyme elevations, reported as adverse events in clinical trials was observed in paediatric patients as compared to adults (14.2% transaminases increased in paediatrics compared to 5.3% in adults). Post-marketing data suggest there might be a higher occurrence of skin reactions (especially erythema) in the paediatric population compared to adults. In the 22 patients less than 2 years old who received voriconazole in a compassionate use programme, the following adverse reactions (for which a relationship to voriconazole could not be excluded) were reported: photosensitivity reaction (1), arrhythmia (1), pancreatitis (1), blood bilirubin increased (1), hepatic enzymes increased (1), rash (1) and papilloedema (1). There have been post-marketing reports of pancreatitis in paediatric patients.

#### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system listed in [Appendix V](#).

#### **4.9 Overdose**

In clinical trials there were 3 cases of accidental overdose. All occurred in paediatric patients, who received up to five times the recommended intravenous dose of voriconazole. A single adverse reaction of photophobia of 10 minutes duration was reported.

There is no known antidote to voriconazole.

Voriconazole is haemodialysed with a clearance of 121 ml/min. In an overdose, haemodialysis may assist in the removal of voriconazole from the body.

## 5. PHARMACOLOGICAL PROPERTIES

### 5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Antimycotics for systemic use, triazole derivatives

ATC code: J02AC03

#### Mechanism of action

Voriconazole is a triazole antifungal agent. The primary mode of action of voriconazole is the inhibition of fungal cytochrome P-450-mediated 14 alpha-lanosterol demethylation, an essential step in fungal ergosterol biosynthesis. The accumulation of 14 alpha-methyl sterols correlates with the subsequent loss of ergosterol in the fungal cell membrane and may be responsible for the antifungal activity of voriconazole. Voriconazole has been shown to be more selective for fungal cytochrome P-450 enzymes than for various mammalian cytochrome P-450 enzyme systems.

#### Pharmacokinetic/Pharmacodynamic Relationship

In 10 therapeutic studies, the median for the average and maximum plasma concentrations in individual subjects across the studies was 2425 ng/ml (inter-quartile range 1193 to 4380 ng/ml) and 3742 ng/ml (inter-quartile range 2027 to 6302 ng/ml), respectively. A positive association between mean, maximum or minimum plasma voriconazole concentration and efficacy in therapeutic studies was not found and this relationship has not been explored in prophylaxis studies.

Pharmacokinetic-Pharmacodynamic analyses of clinical trial data identified positive associations between plasma voriconazole concentrations and both liver function test abnormalities and visual disturbances. Dose adjustments in prophylaxis studies have not been explored.

#### Clinical efficacy and safety

*In vitro*, voriconazole displays broad-spectrum antifungal activity with antifungal potency against *Candida* species (including fluconazole-resistant *C. krusei* and resistant strains of *C. glabrata* and *C. albicans*) and fungicidal activity against all *Aspergillus* species tested. In addition voriconazole shows *in vitro* fungicidal activity against emerging fungal pathogens, including those such as *Scedosporium* or *Fusarium* which have limited susceptibility to existing antifungal agents.

Clinical efficacy defined as partial or complete response, has been demonstrated for *Aspergillus* spp. including *A. flavus*, *A. fumigatus*, *A. terreus*, *A. niger*, *A. nidulans*, *Candida* spp., including *C. albicans*, *C. glabrata*, *C. krusei*, *C. parapsilosis* and *C. tropicalis* and limited numbers of *C. dubliniensis*, *C. inconspicua*, and *C. guilliermondii*, *Scedosporium* spp., including *S. apiospermum*, *S. prolificans* and *Fusarium* spp.

Other treated fungal infections (often with either partial or complete response) included isolated cases of *Alternaria* spp., *Blastomyces dermatitidis*, *Blastoschizomyces capitatus*, *Cladosporium* spp., *Coccidioides immitis*, *Conidiobolus coronatus*, *Cryptococcus neoformans*, *Exserohilum rostratum*, *Exophiala spinifera*, *Fonsecaea pedrosoi*, *Madurella mycetomatis*, *Paecilomyces lilacinus*, *Penicillium* spp. including *P. marneffeii*, *Phialophora richardsiae*, *Scopulariopsis brevicaulis* and *Trichosporon* spp. including *T. beigelii* infections.

*In vitro* activity against clinical isolates has been observed for *Acremonium* spp., *Alternaria* spp., *Bipolaris* spp., *Cladophialophora* spp. and *Histoplasma capsulatum*, with most strains being inhibited by concentrations of voriconazole in the range 0.05 to 2 µg/ml.

*In vitro* activity against the following pathogens has been shown, but the clinical significance is unknown: *Curvularia* spp. and *Sporothrix* spp.

#### Breakpoints

Specimens for fungal culture and other relevant laboratory studies (serology, histopathology) should be obtained prior to therapy to isolate and identify causative organisms. Therapy may be instituted before the results of the cultures and other laboratory studies are known; however, once these results become available, anti-infective therapy should be adjusted accordingly.

The species most frequently involved in causing human infections include *C. albicans*, *C. parapsilosis*, *C. tropicalis*, *C. glabrata* and *C. krusei*, all of which usually exhibit minimal inhibitory concentration (MICs) of less than 1 mg/L for voriconazole.

However, the *in vitro* activity of voriconazole against *Candida* species is not uniform. Specifically, for *C. glabrata*, the MICs of voriconazole for fluconazole-resistant isolates are proportionally higher than are those of fluconazole-susceptible isolates. Therefore, every attempt should be made to identify *Candida* to species level. If antifungal susceptibility testing is available, the MIC results may be interpreted using breakpoint criteria established by European Committee on Antimicrobial Susceptibility Testing (EUCAST).



EUCAST breakpoints

Candida Species	MIC breakpoint (mg/L)	
	≤S (Susceptible)	>R (Resistant)
<i>Candida albicans</i> <sup>1</sup>	0.125	0.125
<i>Candida tropicalis</i> <sup>1</sup>	0.125	0.125
<i>Candida parapsilosis</i> <sup>1</sup>	0.125	0.125
<i>Candida glabrata</i> <sup>2</sup>	Insufficient evidence	
<i>Candida krusei</i> <sup>3</sup>	Insufficient evidence	
Other <i>Candida</i> spp. <sup>4</sup>	Insufficient evidence	
<p><sup>1</sup> Strains with MIC values above the Susceptible (S) breakpoint are rare, or not yet reported. The identification and antimicrobial susceptibility tests on any such isolate must be repeated and if the result is confirmed the isolate sent to a reference laboratory.</p> <p><sup>2</sup> In clinical studies, response to voriconazole in patients with <i>C glabrata</i> infections was 21 % lower compared to <i>C. albicans</i>, <i>C. parapsilosis</i> and <i>C. tropicalis</i>. <i>In vitro</i> data showed a slight increase of resistance of <i>C. glabrata</i> to voriconazole.</p> <p><sup>3</sup> In clinical studies, response to voriconazole in <i>C. krusei</i> infections was similar to <i>C. albicans</i>, <i>C. parapsilosis</i> and <i>C. tropicalis</i>. However, as there were only 9 cases available for EUCAST analysis, there is currently insufficient evidence to set clinical breakpoints for <i>C. krusei</i>.</p> <p><sup>4</sup> EUCAST has not determined non-species related breakpoints for voriconazole.</p>		

Clinical experience

Successful outcome in this section is defined as complete or partial response.

*Aspergillus* infections – efficacy in aspergillosis patients with poor prognosis

Voriconazole has *in vitro* fungicidal activity against *Aspergillus* spp. The efficacy and survival benefit of voriconazole versus conventional amphotericin B in the primary treatment of acute invasive aspergillosis was demonstrated in an open, randomised, multicentre study in 277 immunocompromised patients treated for 12 weeks. Voriconazole was administered intravenously with a loading dose of 6 mg/kg every 12 hours for the first 24 hours followed by a maintenance dose of 4 mg/kg every 12 hours for a minimum of 7 days. Therapy could then be switched to the oral formulation at a dose of 200 mg every 12 hours. Median duration of IV voriconazole therapy was 10 days (range 2-85 days). After IV voriconazole therapy, the median duration of oral voriconazole therapy was 76 days (range 2-232 days).

A satisfactory global response (complete or partial resolution of all attributable symptoms signs, radiographic/bronchoscopic abnormalities present at baseline) was seen in 53 % of voriconazole-treated patients compared to 31 % of patients treated with comparator. The 84-day survival rate for voriconazole was statistically significantly higher than that for the comparator and a clinically and statistically significant benefit was shown in favour of voriconazole for both time to death and time to discontinuation due to toxicity.

This study confirmed findings from an earlier, prospectively designed study where there was a positive outcome in subjects with risk factors for a poor prognosis, including graft versus host disease, and, in particular, cerebral infections (normally associated with almost 100 % mortality).

The studies included cerebral, sinus, pulmonary and disseminated aspergillosis in patients with bone marrow and solid organ transplants, haematological malignancies, cancer and AIDS.

#### Candidaemia in non-neutropenic patients

The efficacy of voriconazole compared to the regimen of amphotericin B followed by fluconazole in the primary treatment of candidaemia was demonstrated in an open, comparative study. Three hundred and seventy non-neutropenic patients (above 12 years of age) with documented candidaemia were included in the study, of whom 248 were treated with voriconazole. Nine subjects in the voriconazole group and 5 in the amphotericin B followed by fluconazole group also had mycologically proven infection in deep tissue. Patients with renal failure were excluded from this study. The median treatment duration was 15 days in both treatment arms. In the primary analysis, successful response as assessed by a Data Review Committee (DRC) blinded to study medicinal product was defined as resolution/improvement in all clinical signs and symptoms of infection with eradication of *Candida* from blood and infected deep tissue sites 12 weeks after the end of therapy (EOT). Patients who did not have an assessment 12 weeks after EOT were counted as failures. In this analysis a successful response was seen in 41 % of patients in both treatment arms.

In a secondary analysis, which utilised DRC assessments at the latest evaluable time point (EOT, or 2, 6, or 12 weeks after EOT) voriconazole and the regimen of amphotericin B followed by fluconazole had successful response rates of 65 % and 71 %, respectively.

The Investigator's assessment of successful outcome at each of these time points is shown in the following table.

<i>Timepoint</i>	<i>Voriconazole</i> (N=248)	<i>Amphotericin B</i> → <i>fluconazole</i> (N=122)
EOT	178 (72 %)	88 (72 %)
2 weeks after EOT	125 (50 %)	62 (51 %)
6 weeks after EOT	104 (42 %)	55 (45 %)
12 weeks after EOT	104 (42 %)	51 (42 %)

#### Serious refractory *Candida* infections

The study comprised 55 patients with serious refractory systemic *Candida* infections (including candidaemia, disseminated and other invasive candidiasis) where prior antifungal treatment, particularly with fluconazole, had been ineffective. Successful response was seen in 24 patients (15 complete, 9 partial responses). In fluconazole-resistant non-*albicans* species, a successful outcome was seen in 3/3 *C. krusei* (complete responses) and 6/8 *C. glabrata* (5 complete, 1 partial response) infections. The clinical efficacy data were supported by limited susceptibility data.

Scedosporium and Fusarium infections

Voriconazole was shown to be effective against the following rare fungal pathogens:

*Scedosporium* spp.: Successful response to voriconazole therapy was seen in 16 (6 complete, 10 partial responses) of 28 patients with *S. apiospermum* and in 2 (both partial responses) of 7 patients with *S. prolificans* infection. In addition, a successful response was seen in 1 of 3 patients with infections caused by more than one organism including *Scedosporium* spp.

*Fusarium* spp.: Seven (3 complete, 4 partial responses) of 17 patients were successfully treated with voriconazole. Of these 7 patients, 3 had eye, 1 had sinus, and 3 had disseminated infection. Four additional patients with fusariosis had an infection caused by several organisms; 2 of them had a successful outcome.

The majority of patients receiving voriconazole treatment of the above mentioned rare infections were intolerant of, or refractory to, prior antifungal therapy.

Primary prophylaxis of invasive fungal infections – efficacy in HSCT recipients without prior proven or probable IFI

Voriconazole was compared to itraconazole as primary prophylaxis in an open-label, comparative, multicenter study of adult and adolescent allogeneic HSCT recipients without prior proven or probable IFI. Success was defined as the ability to continue study drug prophylaxis for 100 days after HSCT (without stopping for >14 days) and survival with no proven or probable IFI for 180 days after HSCT. The modified intent-to-treat (MITT) group included 465 allogeneic HSCT recipients with 45% of patients having AML. From all patients 58% were subject to myeloablative conditions regimens. Prophylaxis with study drug was started immediately after HSCT: 224 received voriconazole and 241 received itraconazole. The median duration of study drug prophylaxis was 96 days for voriconazole and 68 days for itraconazole in the MITT group.

Success rates and other secondary endpoints are presented in the table below:

Study Endpoints	Voriconazole N=224	Itraconazole N=241	Difference in proportions and the 95% confidence interval (CI)	P-Value
Success at day 180*	109 (48.7%)	80 (33.2%)	16.4% (7.7%, 25.1%)**	0.0002**
Success at day 100	121 (54.0%)	96 (39.8%)	15.4% (6.6%, 24.2%)**	0.0006**
Completed at least 100 days of study drug prophylaxis	120 (53.6%)	94 (39.0%)	14.6% (5.6%, 23.5%)	0.0015
Survived to day 180	184 (82.1%)	197 (81.7%)	0.4% (-6.6%, 7.4%)	0.9107
Developed proven or probable IFI to day 180	3 (1.3%)	5 (2.1%)	-0.7% (-3.1%, 1.6%)	0.5390
Developed proven or probable IFI to day 100	2 (0.9%)	4 (1.7%)	-0.8% (-2.8%, 1.3%)	0.4589
Developed proven or probable IFI while on study drug	0	3 (1.2%)	-1.2% (-2.6%, 0.2%)	0.0813

\*Primary endpoint of the study

\*\*Difference in proportions, 95% CI and p-values obtained after adjustment for randomization

The breakthrough IFI rate to Day 180 and the primary endpoint of the study, which is Success at Day 180, for patients with AML and myeloablative conditioning regimens respectively, is presented in the table below:

#### AML

Study endpoints	Voriconazole (N=98)	Itraconazole (N=109)	Difference in proportions and the 95% confidence interval (CI)
Breakthrough IFI – Day 180	1 (1.0%)	2 (1.8%)	-0.8% (-4.0%, 2.4%)**
Success at Day 180*	55 (56.1%)	45 (41.3%)	14.7% (1.7%, 27.7%)***

\*Primary endpoint of study

\*\*Using a margin of 5%, non inferiority is demonstrated

\*\*\*Difference in proportions, 95% CI obtained after adjustment for randomization

#### Myeloablative conditioning regimens

Study endpoints	Voriconazole (N=125)	Itraconazole (N=143)	Difference in proportions and the 95% confidence interval
Breakthrough IFI – Day 180	2 (1.6%)	3 (2.1%)	-0.5% (-3.7%, 2.7%) **
Success at Day 180*	70 (56.0%)	53 (37.1%)	20.1% (8.5%, 31.7%)***

\*Primary endpoint of study

\*\*Using a margin of 5%, non inferiority is demonstrated

\*\*\*Difference in proportions, 95% CI obtained after adjustment for randomization

#### Secondary Prophylaxis of IFI – Efficacy in HSCT recipients with prior proven or probable IFI

Voriconazole was investigated as secondary prophylaxis in an open-label, non-comparative, multicenter study of adult allogeneic HSCT recipients with prior proven or probable IFI. The primary endpoint was the rate of occurrence of proven and probable IFI during the first year after HSCT. The MITT group included 40 patients with prior IFI, including 31 with aspergillosis, 5 with candidiasis, and 4 with other IFI. The median duration of study drug prophylaxis was 95.5 days in the MITT group.

Proven or probable IFIs developed in 7.5% (3/40) of patients during the first year after HSCT, including one candidemia, one scedosporiosis (both relapses of prior IFI), and one zygomycosis. The survival rate at Day 180 was 80.0% (32/40) and at 1 year was 70.0% (28/40).

#### Duration of treatment

In clinical trials, 705 patients received voriconazole therapy for greater than 12 weeks, with 164 patients receiving voriconazole for over 6 months.

### Paediatric population

Fifty-three paediatric patients aged 2 to <18 years were treated with voriconazole in two prospective, open-label, non-comparative, multi-center clinical trials. One study enrolled 31 patients with possible, proven or probable invasive aspergillosis (IA), of whom 14 patients had proven or probable IA and were included in the MITT efficacy analyses. The second study enrolled 22 patients with invasive candidiasis including candidaemia (ICC), and esophageal candidiasis (EC) requiring either primary or salvage therapy, of whom 17 were included in the MITT efficacy analyses. For patients with IA the overall rates of global response at 6 weeks were 64.3% (9/14), the global response rate was 40% (2/5) for patients 2 to <12 years and 77.8% (7/9) for patients 12 to <18 years of age. For patients with ICC the global response rate at EOT was 85.7% (6/7) and for patients with EC the global response rate at EOT was 70% (7/10). The overall rate of response (ICC and EC combined) was 88.9% (8/9) for 2 to <12 years old and 62.5% (5/8) for 12 to <18 years old.

### Clinical studies examining QTc interval

A placebo-controlled, randomized, single-dose, crossover study to evaluate the effect on the QTc interval of healthy volunteers was conducted with three oral doses of voriconazole and ketoconazole. The placebo-adjusted mean maximum increases in QTc from baseline after 800, 1200 and 1600 mg of voriconazole were 5.1, 4.8, and 8.2 msec, respectively and 7.0 msec for ketoconazole 800 mg. No subject in any group had an increase in QTc of  $\geq 60$  msec from baseline. No subject experienced an interval exceeding the potentially clinically relevant threshold of 500 msec.

## **5.2 Pharmacokinetic properties**

### General pharmacokinetic characteristics

The pharmacokinetics of voriconazole have been characterised in healthy subjects, special populations and patients. During oral administration of 200 mg or 300 mg twice daily for 14 days in patients at risk of aspergillosis (mainly patients with malignant neoplasms of lymphatic or haematopoietic tissue), the observed pharmacokinetic characteristics of rapid and consistent absorption, accumulation and non-linear pharmacokinetics were in agreement with those observed in healthy subjects.

The pharmacokinetics of voriconazole are non-linear due to saturation of its metabolism. Greater than proportional increase in exposure is observed with increasing dose. It is estimated that, on average, increasing the oral dose from 200 mg twice daily to 300 mg twice daily leads to a 2.5-fold increase in exposure ( $AUC_{\tau}$ ). The oral maintenance dose of 200 mg (or 100 mg for patients less than 40 kg) achieves a voriconazole exposure similar to 3 mg/kg IV. A 300 mg (or 150 mg for patients less than 40 kg) oral maintenance dose achieves an exposure similar to 4 mg/kg IV. When the recommended intravenous or oral loading dose regimens are administered, plasma concentrations close to steady state are achieved within the first 24 hours of dosing. Without the loading dose, accumulation occurs during twice daily multiple dosing with steady-state plasma voriconazole concentrations being achieved by Day 6 in the majority of subjects.

### Absorption

Voriconazole is rapidly and almost completely absorbed following oral administration, with maximum plasma concentrations ( $C_{max}$ ) achieved 1-2 hours after dosing. The absolute bioavailability of voriconazole after oral administration is estimated to be 96 %. When multiple doses of voriconazole are administered with high fat meals,  $C_{max}$  and  $AUC_{\tau}$  are reduced by 34 % and 24 %, respectively. The absorption of voriconazole is not affected by changes in gastric pH.

#### Distribution

The volume of distribution at steady state for voriconazole is estimated to be 4.6 L/kg, suggesting extensive distribution into tissues. Plasma protein binding is estimated to be 58 %. Cerebrospinal fluid samples from eight patients in a compassionate programme showed detectable voriconazole concentrations in all patients.

#### Biotransformation

*In vitro* studies showed that voriconazole is metabolised by, the hepatic cytochrome P450 isoenzymes, CYP2C19, CYP2C9 and CYP3A4.

The inter-individual variability of voriconazole pharmacokinetics is high.

*In vivo* studies indicated that CYP2C19 is significantly involved in the metabolism of voriconazole. This enzyme exhibits genetic polymorphism. For example, 15-20 % of Asian populations may be expected to be poor metabolisers. For Caucasians and Blacks the prevalence of poor metabolisers is 3-5 %. Studies conducted in Caucasian and Japanese healthy subjects have shown that poor metabolisers have, on average, 4-fold higher voriconazole exposure ( $AUC_{\tau}$ ) than their homozygous extensive metaboliser counterparts. Subjects who are heterozygous extensive metabolisers have on average 2-fold higher voriconazole exposure than their homozygous extensive metaboliser counterparts.

The major metabolite of voriconazole is the N-oxide, which accounts for 72 % of the circulating radiolabelled metabolites in plasma. This metabolite has minimal antifungal activity and does not contribute to the overall efficacy of voriconazole.

#### Elimination

Voriconazole is eliminated via hepatic metabolism with less than 2 % of the dose excreted unchanged in the urine.

After administration of a radiolabelled dose of voriconazole, approximately 80 % of the radioactivity is recovered in the urine after multiple intravenous dosing and 83 % in the urine after multiple oral dosing. The majority (> 94 %) of the total radioactivity is excreted in the first 96 hours after both oral and intravenous dosing.

The terminal half-life of voriconazole depends on dose and is approximately 6 hours at 200 mg (orally). Because of non-linear pharmacokinetics, the terminal half-life is not useful in the prediction of the accumulation or elimination of voriconazole.

#### Pharmacokinetics in special patient groups

##### *Gender*

In an oral multiple dose study,  $C_{max}$  and  $AUC_{\tau}$  for healthy young females were 83 % and 113 % higher, respectively, than in healthy young males (18-45 years). In the same study, no significant differences in  $C_{max}$  and  $AUC_{\tau}$  were observed between healthy elderly males and healthy elderly females ( $\geq 65$  years).

In the clinical programme, no dosage adjustment was made on the basis of gender. The safety profile and plasma concentrations observed in male and female patients were similar. Therefore, no dosage adjustment based on gender is necessary.

#### *Elderly*

In an oral multiple dose study  $C_{max}$  and  $AUC_{\tau}$  in healthy elderly males ( $\geq 65$  years) were 61 % and 86 % higher, respectively, than in healthy young males (18-45 years). No significant differences in  $C_{max}$  and  $AUC_{\tau}$  were observed between healthy elderly females ( $\geq 65$  years) and healthy young females (18-45 years).

In the therapeutic studies no dosage adjustment was made on the basis of age. A relationship between plasma concentrations and age was observed. The safety profile of voriconazole in young and elderly patients was similar and, therefore, no dosage adjustment is necessary for the elderly (see section 4.2).

#### *Paediatric population*

The recommended doses in children and adolescent patients are based on a population pharmacokinetic analysis of data obtained from 112 immunocompromised paediatric patients aged 2 to < 12 years and 26 immunocompromised adolescent patients aged 12 to < 17 years. Multiple intravenous doses of 3, 4, 6, 7 and 8 mg/kg twice daily and multiple oral doses (using the powder for oral suspension) of 4 mg/kg, 6 mg/kg, and 200 mg twice daily were evaluated in 3 paediatric pharmacokinetic studies. Intravenous loading doses of 6 mg/kg IV twice daily on day 1 followed by 4 mg/kg intravenous dose twice daily and 300 mg oral tablets twice daily were evaluated in one adolescent pharmacokinetic study. Larger inter-subject variability was observed in paediatric patients compared to adults.

A comparison of the paediatric and adult population pharmacokinetic data indicated that the predicted total exposure ( $AUC_{\tau}$ ) in children following administration of a 9 mg/kg IV loading dose was comparable to that in adults following a 6 mg/kg IV loading dose. The predicted total exposures in children following IV maintenance doses of 4 and 8 mg/kg twice daily were comparable to those in adults following 3 and 4 mg/kg IV twice daily, respectively. The predicted total exposure in children following an oral maintenance dose of 9 mg/kg (maximum of 350 mg) twice daily was comparable to that in adults following 200 mg oral twice daily. An 8 mg/kg intravenous dose will provide voriconazole exposure approximately 2-fold higher than a 9 mg/kg oral dose.

The higher intravenous maintenance dose in paediatric patients relative to adults reflects the higher elimination capacity in paediatric patients due to a greater liver mass to body mass ratio. Oral bioavailability may, however, be limited in paediatric patients with malabsorption and very low body weight for their age. In that case, intravenous voriconazole administration is recommended.

Voriconazole exposures in the majority of adolescent patients were comparable to those in adults receiving the same dosing regimens. However, lower voriconazole exposure was observed in some young adolescents with low body weight compared to adults. It is likely that these subjects may metabolize voriconazole more similarly to children than to adults. Based on the population pharmacokinetic analysis, 12- to 14-year-old adolescents weighing less than 50 kg should receive children's doses (see section 4.2).

#### Renal impairment

In an oral single-dose (200 mg) study in subjects with normal renal function and mild (creatinine clearance 41-60 ml/min) to severe (creatinine clearance < 20 ml/min) renal impairment, the pharmacokinetics of voriconazole were not significantly affected by renal impairment. The plasma protein binding of voriconazole was similar in subjects with different degrees of renal impairment (see sections 4.2 and 4.4).

#### Hepatic impairment

After an oral single dose (200 mg), AUC was 233 % higher in subjects with mild to moderate hepatic cirrhosis (Child-Pugh A and B) compared with subjects with normal hepatic function. Protein binding of voriconazole was not affected by impaired hepatic function.

In an oral multiple dose study, AUC<sub>τ</sub> was similar in subjects with moderate hepatic cirrhosis (Child-Pugh B) given a maintenance dose of 100 mg twice daily and subjects with normal hepatic function given 200 mg twice daily. No pharmacokinetic data are available for patients with severe hepatic cirrhosis (Child-Pugh C) (see sections 4.2 and 4.4).

### **5.3 Preclinical safety data**

Repeated-dose toxicity studies with voriconazole indicated the liver to be the target organ. Hepatotoxicity occurred at plasma exposures similar to those obtained at therapeutic doses in humans, in common with other antifungal agents. In rats, mice and dogs, voriconazole also induced minimal adrenal changes. Conventional studies of safety pharmacology, genotoxicity or carcinogenic potential did not reveal a special hazard for humans.

In reproduction studies, voriconazole was shown to be teratogenic in rats and embryotoxic in rabbits at systemic exposures equal to those obtained in humans with therapeutic doses. In the pre- and postnatal development study in rats at exposures lower than those obtained in humans with therapeutic doses, voriconazole prolonged the duration of gestation and labour and produced dystocia with consequent maternal mortality and reduced perinatal survival of pups. The effects on parturition are probably mediated by species-specific mechanisms, involving reduction of oestradiol levels, and are consistent with those observed with other azole antifungal agents. Voriconazole administration induced no impairment of male or female fertility in rats at exposures similar to those obtained in humans at therapeutic doses.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

#### Tablet core



Lactose monohydrate  
Pregelatinized maize starch  
Crosscarmellose sodium  
Povidone K-30  
Magnesium stearate

#### Film-coating

Hypromellose 6 mPa\*s  
Titanium dioxide (E171)  
Lactose monohydrate  
Triacetin

### **6.2 Incompatibilities**

Not applicable.

### **6.3 Shelf life**

24 months

### **6.4 Special precautions for storage**

This medicinal product does not require any special storage conditions.

### **6.5 Nature and contents of container**

PVC/Al blisters, paper folding box  
Pack size: 2, 10, 14, 20, 28, 30, 50, 56 or 100 film-coated tablets

Not all pack sizes may be marketed.

### **6.6 Special precautions for disposal**

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

## **7. MARKETING AUTHORISATION HOLDER**

{Name and address}  
<{tel}>  
<{fax}>  
<{e-mail}>  
<[To be completed nationally]>

## **8. MARKETING AUTHORISATION NUMBER(S)**

<[To be completed nationally]>

**9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION**

<[To be completed nationally]>

**10. DATE OF REVISION OF THE TEXT**

<{MM/YYYY}>

<[To be completed nationally]>

## **LABELLING**

### **PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**Paper folding box**

#### **1. NAME OF THE MEDICINAL PRODUCT**

<[Invented name]> 50 mg film-coated tablets  
<[Invented name]> 200 mg film-coated tablets  
Voriconazole

#### **2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each tablet contains 50 mg voriconazole.  
Each tablet contains 200 mg voriconazole.

#### **3. LIST OF EXCIPIENTS**

Contains lactose.  
See leaflet for further information.

#### **4. PHARMACEUTICAL FORM AND CONTENTS**

Film-coated tablet

2 film-coated tablets  
10 film-coated tablets  
14 film-coated tablets  
20 film-coated tablets  
28 film-coated tablets  
30 film-coated tablets  
50 film-coated tablets  
56 film-coated tablets  
100 film-coated tablets

#### **5. METHOD AND ROUTE(S) OF ADMINISTRATION**

Oral use  
Read the package leaflet before use.

#### **6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

DE/H/3920/001-002 Voriconazol 50mg, 200mg FCT\_2018/12 PSUSA – editorial changes and prolonged shelf-life

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

<[To be completed nationally]>

**12. MARKETING AUTHORISATION NUMBER(S)**

<[To be completed nationally]>

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

<[To be completed nationally]>

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

<[Invented name]> 50 mg  
<[Invented name]> 200 mg

DE/H/3920/001-002 Voriconazol 50mg, 200mg FCT\_2018/12 PSUSA – editorial changes and prolonged shelf-life

**17. UNIQUE IDENTIFIER – 2D BARCODE**

2D barcode carrying the unique identifier included.

**18. UNIQUE IDENTIFIER – HUMAN READABLE DATA**

PC:  
SN:  
NN:

DE/H/3920/001-002 Voriconazol 50mg, 200mg FCT\_2018/12 PSUSA – editorial changes and prolonged shelf-life

**MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS**

**Blister**

**1. NAME OF THE MEDICINAL PRODUCT**

<[Invented name]> 50 mg film-coated tablets  
<[Invented name]> 200 mg film-coated tablets  
Voriconazole

**2. NAME OF THE MARKETING AUTHORISATION HOLDER**

<[To be completed nationally]>

**3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Lot

**5. OTHER**

DE/H/3920/001-002 Voriconazol 50mg, 200mg FCT\_2018/12 PSUSA – editorial changes and prolonged shelf-life

## Package leaflet: Information for the patient

<[Invented name]> 50 mg film-coated tablets  
<[Invented name]> 200 mg film-coated tablets

Voriconazole

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor, pharmacist or nurse.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

### What is in this leaflet

1. What <[Invented name]> is and what it is used for
2. What you need to know before you take <[Invented name]>
3. How to take <[Invented name]>
4. Possible side effects
5. How to store <[Invented name]>
6. Contents of the pack and other information

#### 1. What <[Invented name]> is and what it is used for

<[Invented name]> contains the active substance voriconazole. <[Invented name]> is an antifungal medicine. It works by killing or stopping the growth of the fungi that cause infections.

It is used for the treatment of patients (adults and children over the age of 2) with:

- invasive aspergillosis (a type of fungal infection due to *Aspergillus* spp.),
- candidaemia (another type of fungal infection due to *Candida* spp.) in non-neutropenic patients (patients without abnormally low white blood cells count),
- serious invasive *Candida* spp. infections when the fungus is resistant to fluconazole (another antifungal medicine),
- serious fungal infections caused by *Scedosporium* spp. or *Fusarium* spp. (two different species of fungi).

<[Invented name]> is intended for patients with worsening, possibly life-threatening, fungal infections.

Prevention of fungal infections in high risk bone marrow transplant recipients.

This product should only be taken under the supervision of a doctor.

#### 2. What you need to know before you take <[Invented name]>

DE/H/3920/001-002 Voriconazol 50mg, 200mg FCT\_2018/12 PSUSA – editorial changes and prolonged shelf-life

**Do not take <[Invented name]>:**

- if you are allergic to voriconazole or any of the other ingredients of this medicine (listed in section 6).

It is very important that you inform your doctor or pharmacist if you are taking or have taken any other medicines, even those that are obtained without a prescription, or herbal medicines.

The medicines in the following list must not be taken during your course of <[Invented name]> treatment:

- Terfenadine (used for allergy)
- Astemizole (used for allergy)
- Cisapride (used for stomach problems)
- Pimozide (used for treating mental illness)
- Quinidine (used for irregular heart beat)
- Rifampicin (used for treating tuberculosis)
- Efavirenz (used for treating HIV) in doses of 400 mg and above once daily
- Carbamazepine (used to treat seizures)
- Phenobarbital (used for severe insomnia and seizures)
- Ergot alkaloids (e.g., ergotamine, dihydroergotamine; used for migraine)
- Sirolimus (used in transplant patients)
- Ritonavir (used for treating HIV) in doses of 400mg and more twice daily
- St. John's Wort (herbal supplement)

**Warnings and precautions**

Talk to your doctor, pharmacist or nurse before taking <[Invented name]> if:

- you have had an allergic reaction to other azoles.
- you are suffering from, or have ever suffered from liver disease. If you have liver disease, your doctor may prescribe a lower dose of <[Invented name]>. Your doctor should also monitor your liver function while you are being treated with <[Invented name]> by doing blood tests.
- you are known to have cardiomyopathy, irregular heart beat, slow heart rate or an abnormality of electrocardiogram (ECG) called 'long QTc syndrome'.

You should avoid any sunlight and sun exposure while being treated. It is important to cover sun exposed areas of skin and use sunscreen with high sun protection factor (SPF), as an increased sensitivity of skin to the sun's UV rays can occur. These precautions are also applicable to children.

While being treated with <[Invented name]>:

- tell your doctor immediately if you develop
  - sunburn
  - severe skin rash or blisters
  - bone pain.

If you develop skin disorders as described above, your doctor may refer you to a dermatologist, who after consultation may decide that it is important for you to be seen on a regular basis. There is a small chance that skin cancer could develop with long-term use of <[Invented name]>.

Your doctor should monitor the function of your liver and kidney by doing blood tests.

**Children and adolescents**



DE/H/3920/001-002 Voriconazol 50mg, 200mg FCT\_2018/12 PSUSA – editorial changes and prolonged shelf-life  
<[Invented name]> should not be given to children younger than 2 years of age.

### **Other medicines and <[Invented name]>**

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

Some medicines, when taken at the same time as <[Invented name]>, may affect the way <[Invented name]> works or <[Invented name]> may affect the way they work.

Tell your doctor if you are taking the following medicine, as treatment with <[Invented name]> at the same time should be avoided if possible:

- Ritonavir (used for treating HIV) in doses of 100 mg twice daily.

Tell your doctor if you are taking either of the following medicines, as treatment with <[Invented name]> at the same time should be avoided if possible, and a dose adjustment of voriconazole may be required:

- Rifabutin (used for treating tuberculosis). If you are already being treated with rifabutin your blood counts and side effects to rifabutin will need to be monitored.
- Phenytoin (used to treat epilepsy). If you are already being treated with phenytoin your blood concentration of phenytoin will need to be monitored during your treatment with <[Invented name]> and your dose may be adjusted.

Tell your doctor if you are taking any of the following medicines, as a dose adjustment or monitoring may be required to check that the medicines and/ or <[Invented name]> are still having the desired effect:

- Warfarin and other anticoagulants (e.g., phenprocoumon, acenocoumarol; used to slow down clotting of the blood)
- Ciclosporin (used in transplant patients)
- Tacrolimus (used in transplant patients)
- Sulphonylureas (e.g., tolbutamide, glipizide, and glyburide) (used for diabetes)
- Statins (e.g., atorvastatin, simvastatin) (used for lowering cholesterol)
- Benzodiazepines (e.g., midazolam, triazolam) (used for severe insomnia and stress)
- Omeprazole (used for treating ulcers)
- Oral contraceptives (if you take <[Invented name]> whilst using oral contraceptives, you may get side effects such as nausea and menstrual disorders)
- Vinca alkaloids (e.g., vincristine and vinblastine) (used in treating cancer)
- Indinavir and other HIV protease inhibitors (used for treating HIV)
- Non-nucleoside reverse transcriptase inhibitors (e.g., efavirenz, delavirdine, nevirapine) (used for treating HIV) (some doses of efavirenz can NOT be taken at the same time as <[Invented name]>)
- Methadone (used to treat heroin addiction)
- Alfentanil and fentanyl and other short acting opiates such as sufentanil (painkillers used for surgical procedures)
- Oxycodone and other long acting opiates such as hydrocodone (used for moderate to severe pain)
- Non-steroidal anti-inflammatory drugs (e.g., ibuprofen, diclofenac) (used for treating pain and inflammation)
- Fluconazole (used for fungal infections)
- Everolimus (used for treating advanced kidney cancer and in transplant patients)

DE/H/3920/001-002 Voriconazol 50mg, 200mg FCT\_2018/12 PSUSA – editorial changes and prolonged shelf-life

### **Pregnancy and breast-feeding**

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

<[Invented name]> must not be taken during pregnancy, unless indicated by your doctor. Effective contraception must be used in women of childbearing potential. Contact your doctor immediately if you become pregnant while taking <[Invented name]>.

### **Driving and using machines**

<[Invented name]> may cause blurring of vision or uncomfortable sensitivity to light. While affected, do not drive or operate any tools or machines. Contact your doctor if you experience this.

### **<[Invented name]> contains lactose**

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking <[Invented name]>.

## **3. How to take <[Invented name]>**

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

Your doctor will determine your dose depending on your weight and the type of infection you have.

The recommended dose for adults (including elderly patients) is as follows:

	<b>Tablets</b>	
	Patients 40 kg and above	Patients less than 40 kg
<b>Dose for the first 24 hours</b> (Loading Dose)	400 mg every 12 hours for the first 24 hours	200 mg every 12 hours for the first 24 hours
<b>Dose after the first 24 hours</b> (Maintenance Dose)	200 mg twice a day	100 mg twice a day

Depending on your response to treatment, your doctor may increase the daily dose to 300 mg twice a day.

The doctor may decide to decrease the dose if you have mild to moderate cirrhosis.

### **Use in children and adolescents**

The recommended dose for children and teenagers is as follows:

	<b>Tablets</b>	
	Children aged 2 to less than 12 years and teenagers aged 12 to 14 years weighing less than 50 kg	Teenagers aged 12 to 14 years weighing 50 kg or more; and all teenagers older than 14
<b>Dose for the first 24 hours</b> (Loading Dose)	Your treatment will be started as an infusion	400 mg every 12 hours for the first 24 hours

DE/H/3920/001-002 Voriconazol 50mg, 200mg FCT\_2018/12 PSUSA – editorial changes and prolonged shelf-life

<b>Dose after the first 24 hours</b> (Maintenance Dose)	9 mg/kg twice a day (a maximum dose of 350 mg twice daily)	200 mg twice a day
--	--	--------------------

Depending on your response to treatment, your doctor may increase or decrease the daily dose.

Tablets must only be given if the child is able to swallow tablets.

Take your tablet at least one hour before, or one hour after a meal. Swallow the tablet whole with some water.

If you or your child are taking <[Invented name]> for prevention of fungal infections, your doctor may stop giving <[Invented name]> if you or your child develop treatment related side effects.

**If you take more <[Invented name]> than you should**

If you take more tablets than prescribed (or if someone else takes your tablets) you must seek medical advice or go to the nearest hospital casualty department immediately. Take your box of <[Invented name]> tablets with you. You may experience abnormal intolerance to light as a result of taking more <[Invented name]> than you should.

**If you forget to take <[Invented name]>**

It is important to take your <[Invented name]> tablets regularly at the same time each day. If you forget to take one dose, take your next dose when it is due. Do not take a double dose to make up for a forgotten dose.

**If you stop taking <[Invented name]>**

It has been shown that taking all doses at the appropriate times may greatly increase the effectiveness of your medicine. Therefore unless your doctor instructs you to stop treatment, it is important to keep taking <[Invented name]> correctly, as described above.

Continue taking <[Invented name]> until your doctor tells you to stop. Do not stop treatment early because your infection may not be cured. Patients with a weakened immune system or those with difficult infections may require long-term treatment to prevent the infection from returning.

When <[Invented name]> treatment is stopped by your doctor you should not experience any effects.

If you have any further questions on the use of this medicine, ask your doctor, pharmacist or nurse.

**4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

If any side effects occur, most are likely to be minor and temporary. However, some may be serious and need medical attention.

**Serious side effects – Stop taking <[Invented name]> and see a doctor immediately**

- Rash
- Jaundice, changes in blood tests of liver function

DE/H/3920/001-002 Voriconazol 50mg, 200mg FCT\_2018/12 PSUSA – editorial changes and prolonged shelf-life

- Pancreatitis

### **Other side effects**

**Very common:** may affect more than 1 in 10 people

- Visual impairment (change in vision including blurred vision, visual color alterations, abnormal intolerance to visual perception of light, colour blindness, eye disorder, halo vision, night blindness, swinging vision, seeing sparks, visual aura, visual acuity reduced, visual brightness, loss of part of the usual field of vision, spots before the eyes)
- Fever
- Rash
- Nausea, vomiting, diarrhoea
- Headache
- Swelling of the extremities
- Stomach pains
- Breathing difficulties
- Elevated liver enzymes

**Common:** may affect up to 1 in 10 people

- Inflammation of the sinuses, inflammation of the gums, chills, weakness
- Low numbers of some types, including severe, of red (sometimes immune-related) and/or white blood cells (sometimes with fever), low numbers of cells called platelets that help the blood to clot
- 
- Low blood sugar, low blood potassium, low sodium in the blood
- Anxiety, depression, confusion, agitation, inability to sleep, hallucinations
- Seizures, tremors or uncontrolled muscle movements, tingling or abnormal skin sensations, increase in muscle tone, sleepiness, dizziness
- Bleeding in the eye
- Heart rhythm problems including very fast heartbeat, very slow heartbeat, fainting
- Low blood pressure, inflammation of a vein (which may be associated with the formation of a blood clot)
- Acute breathing difficulty, chest pain, swelling of the face (mouth, lips and around eyes), fluid accumulation in the lungs
- Constipation, indigestion, inflammation of the lips
- Jaundice, inflammation of the liver and liver injury
- Skin rashes which may lead to severe blistering and peeling of the skin characterized by a flat, red area on the skin that is covered with small confluent bumps, redness of the skin
- Itchiness
- Hair loss
- Back pain
- Kidney failure, blood in the urine, changes in kidney function tests

**Uncommon:** may affect up to 1 in 100 people

- Flu-like symptoms, irritation and inflammation of the gastrointestinal tract, inflammation of the gastrointestinal tract causing antibiotic associated diarrhoea, inflammation of the lymphatic vessels

DE/H/3920/001-002 Voriconazol 50mg, 200mg FCT\_2018/12 PSUSA – editorial changes and prolonged shelf-life

- Inflammation of the thin tissue that lines the inner wall of the abdomen and covers the abdominal organ
- Enlarged lymph glands (sometimes painful), failure of blood marrow, increased eosinophil
- Allergic reaction or exaggerated immune response
- Depressed function of the adrenal gland, underactive thyroid gland
- Abnormal brain function, Parkinson-like symptoms, nerve injury resulting in numbness, pain, tingling or burning in the hands or feet
- Problems with balance or coordination
- Swelling of the brain
- Double vision, serious conditions of the eye including: pain and inflammation of the eyes and eyelids, abnormal eye movement, damage to the optic nerve resulting in vision impairment, optic disc swelling
- Decreased sensitivity to touch
- Abnormal sense of taste
- Hearing difficulties, ringing in the ears, vertigo
- Inflammation of certain internal organs - pancreas and duodenum, swelling and inflammation of the tongue
- Enlarged liver, liver failure, gallbladder disease, gallstones
- Joint inflammation, inflammation of the veins under the skin (which may be associated with the formation of a blood clot)
- Inflammation of the kidney, proteins in the urine, damage to the kidney
- Very fast heart rate or skipped heartbeats, sometimes with erratic electrical impulses
- Abnormal electrocardiogram (ECG)
- Blood cholesterol increased, blood urea increased
- Allergic skin reactions (sometimes severe), including life-threatening skin condition that causes painful blisters and sores of the skin and mucous membranes, especially in the mouth, inflammation of the skin, hives, sunburn or severe skin reaction following exposure to light or sun, skin redness and irritation, red or purple discoloration of the skin which may be caused by low platelet count, eczema
- Infusion site reaction

**Rare:** may affect up to 1 in 1000 people

- Overactive thyroid gland
- Deterioration of brain function that is a serious complication of liver disease
- Loss of most fibers in the optic nerve, clouding of the cornea, involuntary movement of the eye
- Bullous photosensitivity
- A disorder in which the body's immune system attacks part of the peripheral nervous system
- Heart rhythm or conduction problems (sometimes life threatening)
- Life threatening allergic reaction
- Disorder of blood clotting system
- Allergic skin reactions (sometimes severe), including rapid swelling (oedema) of the dermis, subcutaneous tissue, mucosa and submucosal tissues, itchy or sore patches of thick, red skin with silvery scales of skin, irritation of the skin and mucous membranes, life-threatening skin condition that causes large portions of the epidermis, the skin's outermost layer, to detach from the layers of skin below
- Small dry scaly skin patches, sometimes thick with spikes or 'horns'

DE/H/3920/001-002 Voriconazol 50mg, 200mg FCT\_2018/12 PSUSA – editorial changes and prolonged shelf-life

Side effects with frequency not known:

- Freckles and pigmented spots

Other significant side effects whose frequency is not known, but should be reported to your doctor immediately:

- Skin cancer
- Inflammation of the tissue surrounding the bone
- Red, scaly patches or ring-shaped skin lesions that may be a symptom of an autoimmune disease called cutaneous lupus erythematosus

As <[Invented name]> has been known to affect the liver and the kidney, your doctor should monitor the function of your liver and kidney by doing blood tests. Please advise your doctor if you have any stomach pains or if your stools have a different consistency.

There have been reports of skin cancer in patients treated with voriconazole for long periods of time.

Sunburn or severe skin reaction following exposure to light or sun was experienced more frequently in children. If you or your child develops skin disorders, your doctor may refer you to a dermatologist, who after consultation may decide that it is important for you or your child to be seen on a regular basis. Elevated liver enzymes were also observed more frequently in children.

If any of these side effects persist or are troublesome, please tell your doctor.

### **Reporting of side effects**

If you get any side effects, talk to your doctor or pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via [the national reporting system listed in Appendix V](#). By reporting side effects you can help provide more information on the safety of this medicine.

## **5. How to store <[Invented name]>**

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton/blister after EXP. The expiry date refers to the last day of that month.

This medicine does not require any special storage conditions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

## **6. Contents of the pack and other information**

### **What <[Invented name]> contains**

The active substance is voriconazole. Each tablet contains 50 mg voriconazole.

Each tablet contains 200 mg voriconazole.

DE/H/3920/001-002 Voriconazol 50mg, 200mg FCT\_2018/12 PSUSA – editorial changes and prolonged shelf-life  
The other ingredients are lactose monohydrate, pregelatinized maize starch, croscarmellose sodium, povidone K-30 and magnesium stearate which make up the tablet core and hypromellose 6 mPa\*s, titanium dioxide (E171), lactose monohydrate and triacetin which make up the film-coat.

**What <[Invented name]> looks like and contents of the pack**

<[Invented name]> 50 mg film-coated tablets are white to off- white, round film-coated tablets.

<[Invented name]> 200 mg film-coated tablets are white to off- white, oval film-coated tablets.

Pack size : 2, 10, 14, 20, 28, 30, 50, 56 and 100 film-coated tablets.

Not all pack sizes may be marketed.

**Marketing Authorisation Holder and Manufacturer**

{Name and address}

<{tel}>

<{fax}>

<{e-mail}>

<[To be completed nationally]>

**This medicinal product is authorised in the Member States of the EEA under the following names:**

Germany, Portugal, Spain, Poland, Romania: Voriconazol Zentiva

United Kingdom : Voriconazole Zentiva

**This leaflet was last revised in MM/YYYY.**