

Package leaflet: Information for the user

[Invented name] 2.5 mg film-coated tablets

tadalafil

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What [Invented name] is and what it is used for
2. What you need to know before you take [Invented name]
3. How to take [Invented name]
4. Possible side effects
5. How to store [Invented name]
6. Contents of the pack and other information

1. What [Invented name] is and what it is used for

[Invented name] is a treatment for adult men with erectile dysfunction. This is when a man cannot get, or keep a hard, erect penis suitable for sexual activity. [Invented name] has been shown to significantly improve the ability of obtaining a hard erect penis suitable for sexual activity.

[Invented name] contains the active substance tadalafil which belongs to a group of medicines called phosphodiesterase type 5 inhibitors. Following sexual stimulation [Invented name] works by helping the blood vessels in your penis to relax, allowing the flow of blood into your penis. The result of this is improved erectile function. [Invented name] will not help you if you do not have erectile dysfunction.

It is important to note that [Invented name] does not work if there is no sexual stimulation. You and your partner will need to engage in foreplay, just as you would if you were not taking a medicine for erectile dysfunction.

2. What you need to know before you take [Invented name]

Do NOT take [Invented name] if you:

- are allergic to tadalafil or any of the other ingredients of this medicine (listed in section 6).
- are taking any form of organic nitrate or nitric oxide donors such as amyl nitrite. This is a group of medicines (“nitrates”) used in the treatment of angina pectoris (“chest pain”). [Invented name] has been shown to increase the effects of these medicines. If you are taking any form of nitrate or are unsure tell your doctor (see section “Other medicines and [Invented name]”).
- have serious heart disease or recently had a heart attack within the last 90 days.
- recently had a stroke within the last 6 months.
- have low blood pressure or uncontrolled high blood pressure.
- ever had loss of vision because of non-arteritic anterior ischemic optic neuropathy (NAION), a condition described as “stroke of the eye”.
- are taking riociguat. This drug is used to treat pulmonary arterial hypertension (i.e., high blood pressure in the lungs) and chronic thromboembolic pulmonary hypertension (i.e., high blood pressure in the lungs)

secondary to blood clots). PDE5 inhibitors, such as [Invented name], have been shown to increase the hypotensive effects of this medicine. If you are taking riociguat or are unsure tell your doctor.

Warnings and precautions

Talk to your doctor before taking [Invented name].

Be aware that sexual activity carries a possible risk to patients with heart disease because it puts an extra strain on your heart. If you have a heart problem you should tell your doctor.

Before taking [Invented name], tell your doctor if you have:

- sickle cell anaemia (an abnormality of red blood cells).
- multiple myeloma (cancer of the bone marrow).
- leukaemia (cancer of the blood cells).
- any deformation of your penis.
- a serious liver problem.
- a severe kidney problem.

It is not known if [Invented name] is effective in patients who have had:

- pelvic surgery.
- removal of all or part of the prostate gland in which nerves of the prostate are cut (radical nonnerve-sparing prostatectomy).

If you experience sudden decrease or loss of vision, STOP taking [Invented name] and contact your doctor immediately.

Decreased or sudden hearing loss has been noted in some patients taking tadalafil. Although it is not known if the event is directly related to tadalafil, if you experience decreased or sudden hearing loss, stop taking [Invented name] and contact your doctor immediately.

[Invented name] is not intended for use by women.

Children and adolescents

[Invented name] is not intended for use by children and adolescents under the age of 18.

Other medicines and [Invented name]

Tell your doctor if you are taking, have recently taken or might take any other medicines.

Do NOT take [Invented name] if you are already taking any form of nitrates (see 'Do NOT take [Invented name]').

Some medicines may be affected by [Invented name] or they may affect how well [Invented name] will work. Tell your doctor or pharmacist if you are already taking:

- an alpha blocker (used to treat high blood pressure or urinary symptoms associated with benign prostatic hyperplasia).
- other medicines to treat high blood pressure.
- riociguat.
- a 5-alpha reductase inhibitor (used to treat benign prostatic hyperplasia).
- medicines such as ketoconazole tablets (to treat fungal infections) and protease inhibitors for treatment of AIDS or HIV infection.
- phenobarbital, phenytoin and carbamazepine (anticonvulsant medicines).
- rifampicin, erythromycin, clarithromycin or itraconazole.
- other treatments for erectile dysfunction.

[Invented name] with drink and alcohol

Information on the effect of alcohol is in section 3. Grapefruit juice may affect how well [Invented name] will work and should be taken with caution. Talk to your doctor for further information.

Fertility

When dogs were treated there was reduced sperm development in the testes. A reduction in sperm was seen in some men. These effects are unlikely to lead to a lack of fertility.

Driving and using machines

Some men taking [Invented name] in clinical studies have reported dizziness. Check carefully how you react to the tablets before driving or using machines.

[Invented name] contains lactose and sodium

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.

This medicine contains less than 1 mmol sodium (23 mg) per film-coated tablet, that is to say essentially 'sodium-free'.

3. How to take [Invented name]

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

[Invented name] tablets are for oral use in men only. Swallow the tablet whole with some water. The tablets can be taken with or without food.

The recommended dose is one 5 mg tablet taken once a day at approximately the same time of the day. Your doctor may adjust the dose to 2.5 mg based on your response to [Invented name]. This will be given as a 2.5 mg tablet. Do not take [Invented name] more than once a day.

Once a day dosing of [Invented name] may be useful to men who anticipate having sexual activity two or more times per week. When taken once a day [Invented name] allows you to obtain an erection, when sexually stimulated, at any time point during the 24 hours of the day.

It is important to note that [Invented name] does not work if there is no sexual stimulation. You and your partner will need to engage in foreplay, just as you would if you were not taking a medicine for erectile dysfunction.

Drinking alcohol may affect your ability to get an erection and may temporarily lower your blood pressure. If you have taken or are planning to take [Invented name], avoid excessive drinking (blood alcohol level of 0.08 % or greater), since this may increase the risk of dizziness when standing up.

If you take more [Invented name] than you should

Contact your doctor. You may experience side effects described in section 4.

If you forget to take [Invented name]

Take your dose as soon as you remember but do not take a double dose to make up for a forgotten tablet. You should NOT take [Invented name] more than once a day.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them. These effects are normally mild to moderate in nature.

If you experience any of the following side effects STOP using the medicine and seek medical help immediately:

- allergic reactions including rashes (frequency uncommon).
- chest pain - do not use nitrates but seek immediate medical assistance (frequency uncommon).
- priapism, a prolonged and possibly painful erection after taking [Invented name] (frequency rare). If you have such an erection, which lasts continuously for more than 4 hours you should contact a doctor immediately.
- sudden loss of vision (frequency rare).

Other side effects have been reported:

Common (seen in 1 to 10 in every 100 patients)

- headache
- back pain
- muscle aches
- pain in arms and legs
- facial flushing
- nasal congestion
- indigestion.

Uncommon (seen in 1 to 10 in every 1,000 patients)

- dizziness
- stomach ache
- feeling sick
- being sick (vomiting)
- reflux
- blurred vision
- eye pain
- difficulty in breathing
- presence of blood in urine
- prolonged erection
- pounding heartbeat sensation, a fast heart rate, high blood pressure, low blood pressure
- nose bleeds and ringing in the ears
- swelling of the hands, feet or ankles and feeling tired.

Rare (seen in 1 to 10 in every 10,000 patients)

- fainting
- seizures
- passing memory loss
- swelling of the eyelids, red eyes, sudden decrease or loss of hearing and hives (itchy red welts on the surface of the skin)
- penile bleeding
- presence of blood in semen
- increased sweating.

Heart attack and stroke have also been reported rarely in men taking [Invented name]. Most of these men had known heart problems before taking this medicine.

Partial, temporary, or permanent decrease or loss of vision in one or both eyes has been rarely reported.

Some additional rare side effects have been reported in men taking [Invented name] that were not seen in clinical trials. These include:

- migraine
- swelling of the face
- serious allergic reaction which causes swelling of the face or throat
- serious skin rashes

- some disorders affecting blood flow to the eyes
- irregular heartbeats
- angina
- sudden cardiac death.

The side effect dizziness has been reported more frequently in men over 75 years of age taking [Invented name]. Diarrhoea has been reported more frequently in men over 65 years of age taking [Invented name].

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via [the national reporting system listed in Appendix V](#). By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store [Invented name]

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton and blister after 'EXP'. The expiry date refers to the last day of that month.

This medicinal product does not require any special storage conditions.

Do not throw away any medicines via wastewater <or household waste>. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What [Invented name] contains

- The active substance is tadalafil.
Each 2.5 mg tablet contains 2.5 mg of tadalafil.
- The other ingredients are lactose monohydrate, sodium laurilsulfate, povidone, crospovidone and sodium stearyl fumarate.
- The tablet coating is made of poly(vinyl alcohol), macrogol, titanium dioxide (E171), talc and iron oxide yellow (E172).

What [Invented name] looks like and contents of the pack

[Invented name] 2.5 mg ochre to yellow, round shaped film-coated tablet. On one side debossed with "2.5" and plain on the other side. Diameter: 5.0 mm.

[Invented name] is available in pack sizes of 14, 28, 30 or 56 film-coated tablets in blisters or 28x1 film-coated tablets in perforated unit-dose blisters.

Not all pack sizes may be marketed.

Marketing Authorisation Holder and Manufacturer

[To be completed nationally]

This medicinal product is authorised in the Member States of the EEA under the following names:

This leaflet was last revised in {MM/YYYY}

Package leaflet: Information for the user**[Invented name] 5 mg film-coated tablets**

tadalafil

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What [Invented name] is and what it is used for
2. What you need to know before you take [Invented name]
3. How to take [Invented name]
4. Possible side effects
6. How to store [Invented name]
6. Contents of the pack and other information

1. What [Invented name] is and what it is used for

[Invented name] contains the active substance tadalafil which belongs to a group of medicines called phosphodiesterase type 5 inhibitors.

[Invented name] is used to treat adult men with:

- erectile dysfunction. This is when a man cannot get, or keep a hard, erect penis suitable for sexual activity. [Invented name] has been shown to significantly improve the ability of obtaining a hard erect penis suitable for sexual activity.
Following sexual stimulation [Invented name] works by helping the blood vessels in your penis to relax, allowing the flow of blood into your penis. The result of this is improved erectile function. [Invented name] will not help you if you do not have erectile dysfunction. **It is important to note that [Invented name] for the treatment of erectile dysfunction does not work if there is no sexual stimulation.** You and your partner will need to engage in foreplay, just as you would if you were not taking a medicine for erectile dysfunction.
- urinary symptoms associated with a common condition called benign prostatic hyperplasia. This is when the prostate gland gets bigger with age. Symptoms include difficulty in starting to pass water, a feeling of not completely emptying the bladder and a more frequent need to pass water even at night. [Invented name] improves blood flow to, and relaxes the muscles of, the prostate and bladder which may reduce symptoms of benign prostatic hyperplasia. [Invented name] has been shown to improve these urinary symptoms as early as 1-2 weeks after starting treatment.

2. What you need to know before you take [Invented name]**Do NOT take [Invented name] if you:**

- are allergic to tadalafil or any of the other ingredients of this medicine (listed in section 6).
- are taking any form of organic nitrate or nitric oxide donors such as amyl nitrite. This is a group of medicines (“nitrates”) used in the treatment of angina pectoris (“chest pain”). [Invented name] has been shown to increase the effects of these medicines. If you are taking any form of nitrate or are unsure tell your doctor (see section “Other medicines and [Invented name]”).
- have serious heart disease or recently had a heart attack within the last 90 days.

- recently had a stroke within the last 6 months.
- have low blood pressure or uncontrolled high blood pressure.
- ever had loss of vision because of non-arteritic anterior ischemic optic neuropathy (NAION), a condition described as “stroke of the eye”.
- are taking riociguat. This drug is used to treat pulmonary arterial hypertension (i.e., high blood pressure in the lungs) and chronic thromboembolic pulmonary hypertension (i.e., high blood pressure in the lungs secondary to blood clots). PDE5 inhibitors, such as [Invented name], have been shown to increase the hypotensive effects of this medicine. If you are taking riociguat or are unsure tell your doctor.

Warnings and precautions

Talk to your doctor before taking [Invented name].

Be aware that sexual activity carries a possible risk to patients with heart disease because it puts an extra strain on your heart. If you have a heart problem you should tell your doctor.

Since benign prostatic hyperplasia and prostate cancer may have the same symptoms, your doctor will check you for prostate cancer before starting treatment with [Invented name] for benign prostatic hyperplasia. [Invented name] does NOT treat prostate cancer.

Before taking [Invented name], tell your doctor if you have:

- sickle cell anaemia (an abnormality of red blood cells).
- multiple myeloma (cancer of the bone marrow).
- leukaemia (cancer of the blood cells).
- any deformation of your penis.
- a serious liver problem.
- a severe kidney problem.

It is not known if [Invented name] is effective in patients who have had:

- pelvic surgery.
- removal of all or part of the prostate gland in which nerves of the prostate are cut (radical nonnerve-sparing prostatectomy).

If you experience sudden decrease or loss of vision, STOP taking [Invented name] and contact your doctor immediately.

Decreased or sudden hearing loss has been noted in some patients taking tadalafil. Although it is not known if the event is directly related to tadalafil, if you experience decreased or sudden hearing loss, stop taking [Invented name] and contact your doctor immediately.

[Invented name] is not intended for use by women.

Children and adolescents

[Invented name] is not intended for use by children and adolescents under the age of 18.

Other medicines and [Invented name]

Tell your doctor if you are taking, have recently taken or might take any other medicines.

Do NOT take [Invented name] if you are already taking any form of nitrates (see ‘Do NOT take [Invented name]).

Some medicines may be affected by [Invented name] or they may affect how well [Invented name] will work. Tell your doctor or pharmacist if you are already taking:

- an alpha blocker (used to treat high blood pressure or urinary symptoms associated with benign prostatic hyperplasia).
- other medicines to treat high blood pressure.
- riociguat.

- a 5-alpha reductase inhibitor (used to treat benign prostatic hyperplasia).
- medicines such as ketoconazole tablets (to treat fungal infections) and protease inhibitors for treatment of AIDS or HIV infection.
- phenobarbital, phenytoin and carbamazepine (anticonvulsant medicines).
- rifampicin, erythromycin, clarithromycin or itraconazole.
- other treatments for erectile dysfunction.

[Invented name] with drink and alcohol

Information on the effect of alcohol is in section 3. Grapefruit juice may affect how well [Invented name] will work and should be taken with caution. Talk to your doctor for further information.

Fertility

When dogs were treated there was reduced sperm development in the testes. A reduction in sperm was seen in some men. These effects are unlikely to lead to a lack of fertility.

Driving and using machines

Some men taking [Invented name] in clinical studies have reported dizziness. Check carefully how you react to the tablets before driving or using machines.

[Invented name] contains lactose and sodium

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.

This medicine contains less than 1 mmol sodium (23 mg) per film-coated tablet, that is to say essentially 'sodium-free'.

3. How to take [Invented name]

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

[Invented name] tablets are for oral use in men only. Swallow the tablet whole with some water. The tablets can be taken with or without food.

Drinking alcohol may temporarily lower your blood pressure. If you have taken or are planning to take [Invented name], avoid excessive drinking (blood alcohol level of 0.08% or greater), since this may increase the risk of dizziness when standing up.

For the treatment of erectile dysfunction

The recommended dose is one 5 mg tablet taken once a day at approximately the same time of the day.

Your doctor may adjust the dose to 2.5 mg based on your response to [Invented name]. This will be given as a 2.5 mg tablet. Do not take [Invented name] more than once a day.

Once a day dosing of [Invented name] may be useful to men who anticipate having sexual activity two or more times per week. When taken once a day [Invented name] allows you to obtain an erection, when sexually stimulated, at any time point during the 24 hours of the day.

It is important to note that [Invented name] does not work if there is no sexual stimulation. You and your partner will need to engage in foreplay, just as you would if you were not taking a medicine for erectile dysfunction.

Drinking alcohol may affect your ability to get an erection.

For the treatment of benign prostatic hyperplasia

The dose is one 5 mg tablet taken once a day at approximately the same time of the day.

If you have benign prostatic hyperplasia and erectile dysfunction, the dose remains one 5 mg tablet taken once a day.

Do not take [Invented name] more than once a day.

If you take more [Invented name] than you should

Contact your doctor. You may experience side effects described in section 4.

If you forget to take [Invented name]

Take your dose as soon as you remember but do not take a double dose to make up for a forgotten tablet. You should NOT take [Invented name] more than once a day.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them. These effects are normally mild to moderate in nature.

If you experience any of the following side effects STOP using the medicine and seek medical help immediately:

- allergic reactions including rashes (frequency uncommon).
- chest pain - do not use nitrates but seek immediate medical assistance (frequency uncommon).
- priapism, a prolonged and possibly painful erection after taking [Invented name] (frequency rare). If you have such an erection, which lasts continuously for more than 4 hours you should contact a doctor immediately.
- sudden loss of vision (frequency rare).

Other side effects have been reported:

Common (seen in 1 to 10 in every 100 patients)

- headache
- back pain
- muscle aches
- pain in arms and legs
- facial flushing
- nasal congestion
- indigestion.

Uncommon (seen in 1 to 10 in every 1,000 patients)

- dizziness
- stomach ache
- feeling sick
- being sick (vomiting)
- reflux
- blurred vision
- eye pain
- difficulty in breathing
- presence of blood in urine
- prolonged erection
- pounding heartbeat sensation, a fast heart rate, high blood pressure, low blood pressure
- nose bleeds and ringing in the ears
- swelling of the hands, feet or ankles and feeling tired.

Rare (seen in 1 to 10 in every 10,000 patients)

- fainting
- seizures
- passing memory loss
- swelling of the eyelids, red eyes, sudden decrease or loss of hearing and hives (itchy red welts on the surface of the skin)

- penile bleeding
- presence of blood in semen
- increased sweating.

Heart attack and stroke have also been reported rarely in men taking [Invented name]. Most of these men had known heart problems before taking this medicine.

Partial, temporary, or permanent decrease or loss of vision in one or both eyes has been rarely reported.

Some additional rare side effects have been reported in men taking [Invented name] that were not seen in clinical trials. These include:

- migraine
- swelling of the face
- serious allergic reaction which causes swelling of the face or throat
- serious skin rashes
- some disorders affecting blood flow to the eyes
- irregular heartbeats
- angina
- sudden cardiac death.

The side effect dizziness has been reported more frequently in men over 75 years of age taking [Invented name]. Diarrhoea has been reported more frequently in men over 65 years of age taking [Invented name].

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via **the national reporting system listed in Appendix V**. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store [Invented name]

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton and blister after 'EXP'. The expiry date refers to the last day of that month.

This medicinal product does not require any special storage conditions.

Do not throw away any medicines via wastewater <or household waste>. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What [Invented name] contains

- The active substance is tadalafil.
Each 5 mg tablet contains 5 mg of tadalafil.
- The other ingredients are lactose monohydrate, sodium laurilsulfate, povidone, crospovidone and sodium stearyl fumarate.
- The tablet coating is made of poly(vinyl alcohol), macrogol, titanium dioxide (E171), talc and iron oxide yellow (E172).

What [Invented name] looks like and contents of the pack

[Invented name] 5 mg ochre to yellow, oval shaped film-coated tablet. On one side debossed with "5" and plain on the other side. Length: 8.1 mm, width: 4.1 mm.

[Invented name] is available in pack sizes of 14, 18, 28, 30 or 84 film-coated tablets in blisters or 14 x 1, 28x1, 84 x 1, 98x1 or 112x1 film-coated tablets in perforated unit-dose blisters.

Not all pack sizes may be marketed.

Marketing Authorisation Holder and Manufacturer

[To be completed nationally]

This medicinal product is authorised in the Member States of the EEA under the following names:

This leaflet was last revised in {MM/YYYY}

Package leaflet: Information for the user**[Invented name] 10 mg film-coated tablets**

tadalafil

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What [Invented name] is and what it is used for
2. What you need to know before you take [Invented name]
3. How to take [Invented name]
4. Possible side effects
7. How to store [Invented name]
6. Contents of the pack and other information

1. What [Invented name] is and what it is used for

[Invented name] is a treatment for adult men with erectile dysfunction. This is when a man cannot get, or keep a hard, erect penis suitable for sexual activity. [Invented name] has been shown to significantly improve the ability of obtaining a hard erect penis suitable for sexual activity.

[Invented name] contains the active substance tadalafil which belongs to a group of medicines called phosphodiesterase type 5 inhibitors. Following sexual stimulation [Invented name] works by helping the blood vessels in your penis to relax, allowing the flow of blood into your penis. The result of this is improved erectile function. [Invented name] will not help you if you do not have erectile dysfunction.

It is important to note that [Invented name] does not work if there is no sexual stimulation. You and your partner will need to engage in foreplay, just as you would if you were not taking a medicine for erectile dysfunction.

2. What you need to know before you take [Invented name]**Do NOT take [Invented name] if you:**

- are allergic to tadalafil or any of the other ingredients of this medicine (listed in section 6).
- are taking any form of organic nitrate or nitric oxide donors such as amyl nitrite. This is a group of medicines (“nitrates”) used in the treatment of angina pectoris (“chest pain”). [Invented name] has been shown to increase the effects of these medicines. If you are taking any form of nitrate or are unsure tell your doctor (see section “Other medicines and [Invented name]”).
- have serious heart disease or recently had a heart attack within the last 90 days.
- recently had a stroke within the last 6 months.
- have low blood pressure or uncontrolled high blood pressure.
- ever had loss of vision because of non-arteritic anterior ischemic optic neuropathy (NAION), a condition described as “stroke of the eye”.
- are taking riociguat. This drug is used to treat pulmonary arterial hypertension (i.e., high blood pressure in the lungs) and chronic thromboembolic pulmonary hypertension (i.e., high blood pressure in the lungs)

secondary to blood clots). PDE5 inhibitors, such as [Invented name], have been shown to increase the hypotensive effects of this medicine. If you are taking riociguat or are unsure tell your doctor.

Warnings and precautions

Talk to your doctor before taking [Invented name].

Be aware that sexual activity carries a possible risk to patients with heart disease because it puts an extra strain on your heart. If you have a heart problem you should tell your doctor.

Before taking [Invented name], tell your doctor if you have:

- sickle cell anaemia (an abnormality of red blood cells).
- multiple myeloma (cancer of the bone marrow).
- leukaemia (cancer of the blood cells).
- any deformation of your penis.
- a serious liver problem.
- a severe kidney problem.

It is not known if [Invented name] is effective in patients who have had:

- pelvic surgery.
- removal of all or part of the prostate gland in which nerves of the prostate are cut (radical nonnerve-sparing prostatectomy).

If you experience sudden decrease or loss of vision, STOP taking [Invented name] and contact your doctor immediately.

Decreased or sudden hearing loss has been noted in some patients taking tadalafil. Although it is not known if the event is directly related to tadalafil, if you experience decreased or sudden hearing loss, stop taking [Invented name] and contact your doctor immediately.

[Invented name] is not intended for use by women.

Children and adolescents

[Invented name] is not intended for use by children and adolescents under the age of 18.

Other medicines and [Invented name]

Tell your doctor if you are taking, have recently taken or might take any other medicines.

Do NOT take [Invented name] if you are already taking any form of nitrates (see 'Do NOT take [Invented name]').

Some medicines may be affected by [Invented name] or they may affect how well [Invented name] will work. Tell your doctor or pharmacist if you are already taking:

- an alpha blocker (used to treat high blood pressure or urinary symptoms associated with benign prostatic hyperplasia).
- other medicines to treat high blood pressure.
- riociguat.
- a 5-alpha reductase inhibitor (used to treat benign prostatic hyperplasia).
- medicines such as ketoconazole tablets (to treat fungal infections) and protease inhibitors for treatment of AIDS or HIV infection.
- phenobarbital, phenytoin and carbamazepine (anticonvulsant medicines).
- rifampicin, erythromycin, clarithromycin or itraconazole.
- other treatments for erectile dysfunction.

[Invented name] with drink and alcohol

Information on the effect of alcohol is in section 3. Grapefruit juice may affect how well [Invented name] will work and should be taken with caution. Talk to your doctor for further information.

Fertility

When dogs were treated there was reduced sperm development in the testes. A reduction in sperm was seen in some men. These effects are unlikely to lead to a lack of fertility.

Driving and using machines

Some men taking [Invented name] in clinical studies have reported dizziness. Check carefully how you react to the tablets before driving or using machines.

[Invented name] contains lactose and sodium

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.

This medicine contains less than 1 mmol sodium (23 mg) per film-coated tablet, that is to say essentially 'sodium-free'.

3. How to take [Invented name]

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

[Invented name] tablets are for oral use in men only. Swallow the tablet whole with some water. The tablets can be taken with or without food.

The recommended starting dose is one 10 mg tablet before sexual activity. If the effect of this dose is too weak your doctor may increase the dose to 20 mg. [Invented name] tablets are for oral use.

You may take a [Invented name] tablet at least 30 minutes before sexual activity.

[Invented name] may still be effective up to 36 hours after taking the tablet.

Do not take [Invented name] more than once a day. [Invented name] 10 mg and 20 mg is intended for use prior to anticipated sexual activity and is not recommended for continuous daily use.

It is important to note that [Invented name] does not work if there is no sexual stimulation. You and your partner will need to engage in foreplay, just as you would if you were not taking a medicine for erectile dysfunction.

Drinking alcohol may affect your ability to get an erection and may temporarily lower your blood pressure. If you have taken or are planning to take [Invented name], avoid excessive drinking (blood alcohol level of 0.08 % or greater), since this may increase the risk of dizziness when standing up.

If you take more [Invented name] than you should

Contact your doctor. You may experience side effects described in section 4.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them. These effects are normally mild to moderate in nature.

If you experience any of the following side effects STOP using the medicine and seek medical help immediately:

- allergic reactions including rashes (frequency uncommon).
- chest pain - do not use nitrates but seek immediate medical assistance (frequency uncommon).

- priapism, a prolonged and possibly painful erection after taking [Invented name] (frequency rare). If you have such an erection, which lasts continuously for more than 4 hours you should contact a doctor immediately.
- sudden loss of vision (frequency rare).

Other side effects have been reported:

Common (seen in 1 to 10 in every 100 patients)

- headache
- back pain
- muscle aches
- pain in arms and legs
- facial flushing
- nasal congestion
- indigestion.

Uncommon (seen in 1 to 10 in every 1,000 patients)

- dizziness
- stomach ache
- feeling sick
- being sick (vomiting)
- reflux
- blurred vision
- eye pain
- difficulty in breathing
- presence of blood in urine
- prolonged erection
- pounding heartbeat sensation, a fast heart rate, high blood pressure, low blood pressure
- nose bleeds and ringing in the ears
- swelling of the hands, feet or ankles and feeling tired.

Rare (seen in 1 to 10 in every 10,000 patients)

- fainting
- seizures
- passing memory loss
- swelling of the eyelids, red eyes, sudden decrease or loss of hearing and hives (itchy red welts on the surface of the skin)
- penile bleeding
- presence of blood in semen
- increased sweating.

Heart attack and stroke have also been reported rarely in men taking [Invented name]. Most of these men had known heart problems before taking this medicine.

Partial, temporary, or permanent decrease or loss of vision in one or both eyes has been rarely reported.

Some additional rare side effects have been reported in men taking [Invented name] that were not seen in clinical trials. These include:

- migraine
- swelling of the face
- serious allergic reaction which causes swelling of the face or throat
- serious skin rashes
- some disorders affecting blood flow to the eyes
- irregular heartbeats
- angina
- sudden cardiac death.

The side effect dizziness has been reported more frequently in men over 75 years of age taking [Invented name]. Diarrhoea has been reported more frequently in men over 65 years of age taking [Invented name].

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the national reporting system listed in Appendix V. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store [Invented name]

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton and blister after 'EXP'. The expiry date refers to the last day of that month.

This medicinal product does not require any special storage conditions.

Do not throw away any medicines via wastewater <or household waste>. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What [Invented name] contains

- The active substance is tadalafil.
Each 10 mg tablet contains 10 mg of tadalafil.
- The other ingredients are lactose monohydrate, sodium laurilsulfate, povidone, crospovidone and sodium stearyl fumarate.
- The tablet coating is made of poly(vinyl alcohol), macrogol, titanium dioxide (E171), talc and iron oxide yellow (E172).

What [Invented name] looks like and contents of the pack

[Invented name] 10 mg ochre to yellow, oval shaped film-coated tablet. Debossed with "10" on one side. The other side of the tablet is plain. Length: 11.1 mm, width: 6.4 mm.

[Invented name] is available in pack sizes of 4, 12, 18 or 28 film-coated tablets in blisters or 4x1 or 24x1 film-coated tablets in perforated unit-dose blisters.

Not all pack sizes may be marketed.

Marketing Authorisation Holder and Manufacturer

[To be completed nationally]

This medicinal product is authorised in the Member States of the EEA under the following names:

This leaflet was last revised in {MM/YYYY}

Package leaflet: Information for the user

[Invented name] 20 mg film-coated tablets

tadalafil

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet

1. What [Invented name] is and what it is used for
2. What you need to know before you take [Invented name]
3. How to take [Invented name]
4. Possible side effects
8. How to store [Invented name]
6. Contents of the pack and other information

1. What [Invented name] is and what it is used for

[Invented name] is a treatment for adult men with erectile dysfunction. This is when a man cannot get, or keep a hard, erect penis suitable for sexual activity. [Invented name] has been shown to significantly improve the ability of obtaining a hard erect penis suitable for sexual activity.

[Invented name] contains the active substance tadalafil which belongs to a group of medicines called phosphodiesterase type 5 inhibitors. Following sexual stimulation [Invented name] works by helping the blood vessels in your penis to relax, allowing the flow of blood into your penis. The result of this is improved erectile function. [Invented name] will not help you if you do not have erectile dysfunction.

It is important to note that [Invented name] does not work if there is no sexual stimulation. You and your partner will need to engage in foreplay, just as you would if you were not taking a medicine for erectile dysfunction.

2. What you need to know before you take [Invented name]

Do NOT take [Invented name] if you:

- are allergic to tadalafil or any of the other ingredients of this medicine (listed in section 6).
- are taking any form of organic nitrate or nitric oxide donors such as amyl nitrite. This is a group of medicines (“nitrates”) used in the treatment of angina pectoris (“chest pain”). [Invented name] has been shown to increase the effects of these medicines. If you are taking any form of nitrate or are unsure tell your doctor (see section “Other medicines and [Invented name]”).
- have serious heart disease or recently had a heart attack within the last 90 days.
- recently had a stroke within the last 6 months.
- have low blood pressure or uncontrolled high blood pressure.
- ever had loss of vision because of non-arteritic anterior ischemic optic neuropathy (NAION), a condition described as “stroke of the eye”.
- are taking riociguat. This drug is used to treat pulmonary arterial hypertension (i.e., high blood pressure in the lungs) and chronic thromboembolic pulmonary hypertension (i.e., high blood pressure in the lungs)

secondary to blood clots). PDE5 inhibitors, such as [Invented name], have been shown to increase the hypotensive effects of this medicine. If you are taking riociguat or are unsure tell your doctor.

Warnings and precautions

Talk to your doctor before taking [Invented name].

Be aware that sexual activity carries a possible risk to patients with heart disease because it puts an extra strain on your heart. If you have a heart problem you should tell your doctor.

Before taking [Invented name], tell your doctor if you have:

- sickle cell anaemia (an abnormality of red blood cells).
- multiple myeloma (cancer of the bone marrow).
- leukaemia (cancer of the blood cells).
- any deformation of your penis.
- a serious liver problem.
- a severe kidney problem.

It is not known if [Invented name] is effective in patients who have had:

- pelvic surgery.
- removal of all or part of the prostate gland in which nerves of the prostate are cut (radical nonnerve-sparing prostatectomy).

If you experience sudden decrease or loss of vision, STOP taking [Invented name] and contact your doctor immediately.

Decreased or sudden hearing loss has been noted in some patients taking tadalafil. Although it is not known if the event is directly related to tadalafil, if you experience decreased or sudden hearing loss, stop taking [Invented name] and contact your doctor immediately.

[Invented name] is not intended for use by women.

Children and adolescents

[Invented name] is not intended for use by children and adolescents under the age of 18.

Other medicines and [Invented name]

Tell your doctor if you are taking, have recently taken or might take any other medicines.

Do NOT take [Invented name] if you are already taking any form of nitrates (see 'Do NOT take [Invented name]').

Some medicines may be affected by [Invented name] or they may affect how well [Invented name] will work. Tell your doctor or pharmacist if you are already taking:

- an alpha blocker (used to treat high blood pressure or urinary symptoms associated with benign prostatic hyperplasia).
- other medicines to treat high blood pressure.
- riociguat.
- a 5-alpha reductase inhibitor (used to treat benign prostatic hyperplasia).
- medicines such as ketoconazole tablets (to treat fungal infections) and protease inhibitors for treatment of AIDS or HIV infection.
- phenobarbital, phenytoin and carbamazepine (anticonvulsant medicines).
- rifampicin, erythromycin, clarithromycin or itraconazole.
- other treatments for erectile dysfunction.

[Invented name] with drink and alcohol

Information on the effect of alcohol is in section 3. Grapefruit juice may affect how well [Invented name] will work and should be taken with caution. Talk to your doctor for further information.

Fertility

When dogs were treated there was reduced sperm development in the testes. A reduction in sperm was seen in some men. These effects are unlikely to lead to a lack of fertility.

Driving and using machines

Some men taking [Invented name] in clinical studies have reported dizziness. Check carefully how you react to the tablets before driving or using machines.

[Invented name] contains lactose and sodium

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.

This medicine contains less than 1 mmol sodium (23 mg) per film-coated tablet, that is to say essentially 'sodium-free'.

3. How to take [Invented name]

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

[Invented name] tablets are for oral use in men only.

Once-a-day dosing

The recommended dose is **5 mg** (corresponds to a **quarter of a tablet**) taken once a day at approximately the same time of the day. Your doctor may adjust the dose to 2.5 mg based on your response to [Invented name]. This will be given as a 2.5 mg tablet. Do not take [Invented name] more than once a day.

Once-a-day dosing of [Invented name] may be useful to men who anticipate having sexual activity two or more times per week. When taken once a day [Invented name] allows you to obtain an erection, when sexually stimulated, at any time point during the 24 hours of the day.

On-demand dosing

The recommended starting dose is **10 mg** (corresponds to a **half a tablet**) before sexual activity. If the effect of this dose is too weak your doctor may increase the dose to **20 mg** (corresponds to **one tablet**).

You may take the dose at least 30 minutes before sexual activity.

[Invented name] may still be effective up to 36 hours after taking the dose.

Do not take doses of 10 mg and 20 mg more than once a day. They are intended for use prior to anticipated sexual activity and is not recommended for continuous daily use.

It is important to note that [Invented name] does not work if there is no sexual stimulation. You and your partner will need to engage in foreplay, just as you would if you were not taking a medicine for erectile dysfunction.

Drinking alcohol may affect your ability to get an erection and may temporarily lower your blood pressure. If you have taken or are planning to take [Invented name], avoid excessive drinking (blood alcohol level of 0.08% or greater), since this may increase the risk of dizziness when standing up.


Method of administration

Depending on your dosage regimen (*once-a-day* or *on-demand*) the tablets can be divided in quarters (*once-a-day*) or halves (*on-demand*).

[Invented name] is for oral use and can be taken with or without food.

Swallow the tablet or part of the tablet with some water.

Dividing the tablet into quarters (with 5 mg tadalafil for *once-a-day* dosing)

Place the tablet with the score lines uppermost on a hard, level surface (e.g. tabletop).	
Press down forcefully for a short time with your thumb (or index finger) onto the scored upper side of the tablet until it breaks into four equal parts.	

Dividing the tablet into halves (with 10 mg tadalafil for *on-demand* dosing)

Place the tablet with the score lines uppermost on a hard, level surface (e.g. tabletop).	
Place the thumbs (or index fingers) of both hands at the same time on the left and right side of the shorter score line of the tablet. Press down forcefully for a short time with both thumbs (or index fingers) on the left and right side of the outside of the tablet until it breaks into two equal parts.	

If you take more [Invented name] than you should

Contact your doctor. You may experience side effects described in section 4.

If you forget to take [Invented name]

If you take [Invented name] once a day take your dose as soon as you remember but do not take a double dose to make up for a forgotten tablet. You should NOT take [Invented name] more than once a day.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them. These effects are normally mild to moderate in nature.

If you experience any of the following side effects STOP using the medicine and seek medical help immediately:

- allergic reactions including rashes (frequency uncommon).
- chest pain - do not use nitrates but seek immediate medical assistance (frequency uncommon).
- priapism, a prolonged and possibly painful erection after taking [Invented name] (frequency rare). If you have such an erection, which lasts continuously for more than 4 hours you should contact a doctor immediately.
- sudden loss of vision (frequency rare).

Other side effects have been reported:

Common (seen in 1 to 10 in every 100 patients)

- headache
- back pain
- muscle aches
- pain in arms and legs
- facial flushing
- nasal congestion
- indigestion.

Uncommon (seen in 1 to 10 in every 1,000 patients)

- dizziness
- stomach ache
- feeling sick
- being sick (vomiting)
- reflux
- blurred vision
- eye pain
- difficulty in breathing
- presence of blood in urine
- prolonged erection
- pounding heartbeat sensation, a fast heart rate, high blood pressure, low blood pressure
- nose bleeds and ringing in the ears
- swelling of the hands, feet or ankles and feeling tired.

Rare (seen in 1 to 10 in every 10,000 patients)

- fainting
- seizures
- passing memory loss
- swelling of the eyelids, red eyes, sudden decrease or loss of hearing and hives (itchy red welts on the surface of the skin)
- penile bleeding
- presence of blood in semen
- increased sweating.

Heart attack and stroke have also been reported rarely in men taking [Invented name]. Most of these men had known heart problems before taking this medicine.

Partial, temporary, or permanent decrease or loss of vision in one or both eyes has been rarely reported.

Some additional rare side effects have been reported in men taking [Invented name] that were not seen in clinical trials. These include:

- migraine
- swelling of the face
- serious allergic reaction which causes swelling of the face or throat
- serious skin rashes
- some disorders affecting blood flow to the eyes
- irregular heartbeats
- angina
- sudden cardiac death.

The side effect dizziness has been reported more frequently in men over 75 years of age taking [Invented name]. Diarrhoea has been reported more frequently in men over 65 years of age taking [Invented name].

Reporting of side effects

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the national reporting system listed in Appendix V. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store [Invented name]

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton and blister after 'EXP'. The expiry date refers to the last day of that month.

This medicinal product does not require any special storage conditions.

Do not throw away any medicines via wastewater <or household waste>. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What [Invented name] contains

- The active substance is tadalafil.
Each 20 mg tablet contains 20 mg of tadalafil.
- The other ingredients are lactose monohydrate, sodium laurilsulfate, povidone, crospovidone and sodium stearyl fumarate.
- The tablet coating is made of poly(vinyl alcohol), macrogol, titanium dioxide (E171), talc and iron oxide yellow (E172).

What [Invented name] looks like and contents of the pack

[Invented name] 20 mg ochre to yellow, oval shaped film-coated tablet. One side of the tablet is bi-scored. The other side of the tablet is debossed with "20". Length: 15.0 mm, width: 9.0 mm.
The tablet can be divided into two or four equal doses.

[Invented name] is available in pack sizes of 2, 4, 5, 8, 10, 12, 15, 18, 24, 28, 36, 48 or 56 film-coated tablets in blisters or 2x1, 4x1, 8x1, 12x1, 24x1 or 36x1 film-coated tablets in perforated unit-dose blisters.

Not all pack sizes may be marketed.

Marketing Authorisation Holder and Manufacturer

[To be completed nationally]

This medicinal product is authorised in the Member States of the EEA under the following names:

This leaflet was last revised in {MM/YYYY}