

## SUMMARY OF PRODUCT CHARACTERISTICS

### 1. NAME OF THE MEDICINAL PRODUCT

<Product name> 25 mg film-coated tablets  
<Product name> 50 mg film-coated tablets  
<Product name> 100 mg film-coated tablets  
<Product name> 150 mg film-coated tablets

### 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

<Product name> 25 mg film-coated tablets  
Each film-coated tablet contains 25 mg erlotinib (as erlotinib hydrochloride).  
<Product name> 50 mg film-coated tablets  
Each film-coated tablet contains 50 mg erlotinib (as erlotinib hydrochloride).  
<Product name> 100 mg film-coated tablets  
Each film-coated tablet contains 100 mg erlotinib (as erlotinib hydrochloride).  
<Product name> 150 mg film-coated tablets  
Each film-coated tablet contains 150 mg erlotinib (as erlotinib hydrochloride).

#### Excipient(s) with known effect:

<Product name> 25 mg film-coated tablets  
Each film-coated tablet contains 23.98 mg Lactose monohydrate.  
<Product name> 50 mg film-coated tablets  
Each film-coated tablet contains 47.97 mg Lactose monohydrate.  
<Product name> 100 mg film-coated tablets  
Each film-coated tablet contains 95.93 mg Lactose monohydrate.  
1<Product name> 150 mg film-coated tablets  
Each film-coated tablet contains 143.90 mg Lactose monohydrate.

For the full list of excipients, see section 6.1.

### 3. PHARMACEUTICAL FORM

Film-coated tablet.

<Product name> 25 mg film-coated tablets  
White to yellowish, round biconvex, film-coated tablet, engraved with “25” on one side. The diameter of the tablet is 6.1 mm ± 5 %.  
<Product name> 50 mg film-coated tablets  
White to yellowish, round biconvex, film-coated tablet, engraved with “50” on one side. The diameter of the tablet is 7.6 mm ± 5 %.  
<Product name> 100 mg film-coated tablets  
White to yellowish, round biconvex, film-coated tablet, engraved with “100” on one side. The diameter of the tablet is 8.9 mm ± 5 %.  
<Product name> 150 mg film-coated tablets  
White to yellowish, round biconvex, film-coated tablet, engraved with “150” on one side. The diameter of the tablet is 10.5 mm ± 5 %.

### 4. CLINICAL PARTICULARS

## 4.1 Therapeutic indications

### Non-Small Cell Lung Cancer (NSCLC):

<Product name> is indicated for the first-line treatment of patients with locally advanced or metastatic non-small cell lung cancer (NSCLC) with EGFR activating mutations.

<Product name> is also indicated for switch maintenance treatment in patients with locally advanced or metastatic NSCLC with EGFR activating mutations and stable disease after first-line chemotherapy.

<Product name> is also indicated for the treatment of patients with locally advanced or metastatic NSCLC after failure of at least one prior chemotherapy regimen.

When prescribing <Product name>, factors associated with prolonged survival should be taken into account.

No survival benefit or other clinically relevant effects of the treatment have been demonstrated in patients with Epidermal Growth Factor Receptor (EGFR)-IHC negative tumours (see section 5.1).

### Pancreatic cancer:

<Product name> in combination with gemcitabine is indicated for the treatment of patients with metastatic pancreatic cancer.

When prescribing <Product name>, factors associated with prolonged survival should be taken into account (see sections 4.2 and 5.1).

No survival advantage could be shown for patients with locally advanced disease.

## 4.2 Posology and method of administration

<Product name> treatment should be supervised by a physician experienced in the use of anti-cancer therapies.

### Posology

#### Patients with Non-Small Cell Lung Cancer:

EGFR mutation testing should be performed prior to initiation of <Product name> therapy in chemo-naïve patients with advanced or metastatic NSCLC.

The recommended daily dose of <Product name> is 150 mg taken at least one hour before or two hours after the ingestion of food.

#### Patients with pancreatic cancer:

The recommended daily dose of <Product name> is 100 mg taken at least one hour before or two hours after the ingestion of food, in combination with gemcitabine (see the summary of product characteristics of gemcitabine for the pancreatic cancer indication). In patients who do not develop rash within the first 4 – 8 weeks of treatment, further <Product name> treatment should be re-assessed (see section 5.1).

When dose adjustment is necessary, the dose should be reduced in 50 mg steps (see section 4.4).

For MSs where 25 mg, 50 mg and 100 mg are not registered:

<Product name> is not available in 25 mg, 50 mg and 100 mg strengths. For these dosages, you should take other medicinal products available on the market.

<Product name> is available in strengths of 25 mg, 50 mg, 100 mg and 150 mg.

Concomitant use of CYP3A4 substrates and modulators may require dose adjustment (see section 4.5).

Patients with hepatic impairment: Erlotinib is eliminated by hepatic metabolism and biliary excretion. Although erlotinib exposure was similar in patients with moderately impaired hepatic function (Child- Pugh score 7-9) compared with patients with adequate hepatic function, caution should be used when administering <Product name> to patients with hepatic impairment. Dose reduction or interruption of <Product name> should be considered if severe adverse reactions occur. The safety and efficacy of erlotinib has not been studied in patients with severe hepatic dysfunction (AST/SGOT and ALT/SGPT > 5 x ULN). Use of <Product name> in patients with severe hepatic dysfunction is not recommended (see section 5.2).

Patients with renal impairment: The safety and efficacy of erlotinib has not been studied in patients with renal impairment (serum creatinine concentration >1.5 times the upper normal limit). Based on pharmacokinetic data no dose adjustments appear necessary in patients with mild or moderate renal impairment (see section 5.2). Use of <Product name> in patients with severe renal impairment is not recommended.

#### *Paediatric population*

The safety and efficacy of erlotinib in patients under the age of 18 years has not been established. Use of <Product name> in paediatric patients is not recommended.

Smokers: Cigarette smoking has been shown to reduce erlotinib exposure by 50-60%. The maximum tolerated dose of <Product name> in NSCLC patients who currently smoke cigarettes was 300 mg. Efficacy and long term safety of a dose higher than the recommended starting doses have not been established in patients who continue to smoke cigarettes (see sections 4.5 and 5.2). Therefore, current smokers should be advised to stop smoking, as plasma concentrations of erlotinib in smokers as compared to non-smokers are reduced.

#### Method of administration

For oral use.

### **4.3 Contraindications**

Hypersensitivity to erlotinib or to any of the excipients listed in section 6.1.

### **4.4 Special warnings and precautions for use**

#### Assessment of EGFR mutation status

When assessing the EGFR mutation status of a patient, it is important that a well-validated and robust methodology is chosen to avoid false negative or false positive determinations.

#### Smokers

Current smokers should be advised to stop smoking, as plasma concentrations of erlotinib in smokers as compared to non-smokers are reduced. The degree of reduction is likely to be clinically significant (see section 4.5).

#### Interstitial Lung Disease

Cases of interstitial lung disease (ILD)-like events, including fatalities, have been reported uncommonly in patients receiving erlotinib for treatment of non-small cell lung cancer (NSCLC), pancreatic cancer or other advanced solid tumours. In the pivotal study BR.21 in NSCLC, the incidence of ILD (0.8%) was the same in both the placebo and erlotinib groups. In a meta-analysis of NSCLC randomized controlled clinical trials (excluding phase I and single-arm phase II studies due to lack of control groups), the incidence of ILD-like events

was 0.9% on erlotinib compared to 0.4% in patients in the control arms. In the pancreatic cancer study in combination with gemcitabine, the incidence of ILD-like events was 2.5% in the erlotinib plus gemcitabine group versus 0.4% in the placebo plus gemcitabine treated group. Reported diagnoses in patients suspected of having ILD-like events included pneumonitis, radiation pneumonitis, hypersensitivity pneumonitis, interstitial pneumonia, interstitial lung disease, obliterative bronchiolitis, pulmonary fibrosis, Acute Respiratory Distress Syndrome (ARDS), alveolitis, and lung infiltration. Symptoms started from a few days to several months after initiating erlotinib therapy. Confounding or contributing factors such as concomitant or prior chemotherapy, prior radiotherapy, pre-existing parenchymal lung disease, metastatic lung disease, or pulmonary infections were frequent. A higher incidence of ILD (approximately 5% with a mortality rate of 1.5%) is seen among patients in studies conducted in Japan.

In patients who develop acute onset of new and/or progressive unexplained pulmonary symptoms such as dyspnoea, cough and fever, erlotinib therapy should be interrupted pending diagnostic evaluation. Patients treated concurrently with erlotinib and gemcitabine should be monitored carefully for the possibility to develop ILD-like toxicity. If ILD is diagnosed, erlotinib should be discontinued and appropriate treatment initiated as necessary (see section 4.8).

#### Diarrhoea, dehydration, electrolyte imbalance and renal failure

Diarrhoea (including very rare cases with a fatal outcome) has occurred in approximately 50% of patients on erlotinib and moderate or severe diarrhoea should be treated with e.g. loperamide. In some cases dose reduction may be necessary. In the clinical studies doses were reduced by 50 mg steps. Dose reductions by 25 mg steps have not been investigated. In the event of severe or persistent diarrhoea, nausea, anorexia, or vomiting associated with dehydration, erlotinib therapy should be interrupted and appropriate measures should be taken to treat the dehydration (see section 4.8). There have been rare reports of hypokalaemia and renal failure (including fatalities). Some cases were secondary to severe dehydration due to diarrhoea, vomiting and/or anorexia, while others were confounded by concomitant chemotherapy. In more severe or persistent cases of diarrhoea, or cases leading to dehydration, particularly in groups of patients with aggravating risk factors (especially concomitant chemotherapy and other medications, symptoms or diseases or other predisposing conditions including advanced age), erlotinib therapy should be interrupted and appropriate measures should be taken to intensively rehydrate the patients intravenously. In addition, renal function and serum electrolytes including potassium should be monitored in patients at risk of dehydration.

#### Hepatitis, hepatic failure

Rare cases of hepatic failure (including fatalities) have been reported during use of erlotinib. Confounding factors have included pre-existing liver disease or concomitant hepatotoxic medications. Therefore, in such patients, periodic liver function testing should be considered. erlotinib dosing should be interrupted if changes in liver function are severe (see section 4.8). <Product name> is not recommended for use in patients with severe hepatic dysfunction.

#### Gastrointestinal perforation

Patients receiving <Product name> are at increased risk of developing gastrointestinal perforation, which was observed uncommonly (including some cases with a fatal outcome). Patients receiving concomitant anti-angiogenic agents, corticosteroids, NSAIDs, and/or taxane based chemotherapy, or who have prior history of peptic ulceration or diverticular disease are at increased risk. <Product name> should be permanently discontinued in patients who develop gastrointestinal perforation (see section 4.8).

#### Bullous and exfoliative skin disorders

Bullous, blistering and exfoliative skin conditions have been reported, including very rare

cases suggestive of Stevens-Johnson syndrome/Toxic epidermal necrolysis, which in some cases were fatal (see section 4.8). <Product name> treatment should be interrupted or discontinued if the patient develops severe bullous, blistering or exfoliating conditions. Patients with bullous and exfoliative skin disorders should be tested for skin infection and treated according to local management guidelines.

#### Ocular disorders

Patients presenting with signs and symptoms suggestive of keratitis such as acute or worsening: eye inflammation, lacrimation, light sensitivity, blurred vision, eye pain and/or red eye should be referred promptly to an ophthalmology specialist. If a diagnosis of ulcerative keratitis is confirmed, treatment with <Product name> should be interrupted or discontinued. If keratitis is diagnosed, the benefits and risks of continuing treatment should be carefully considered. <Product name> should be used with caution in patients with a history of keratitis, ulcerative keratitis or severe dry eye. Contact lens use is also a risk factor for keratitis and ulceration. Very rare cases of corneal perforation or ulceration have been reported during use of erlotinib (see section 4.8).

#### Interactions with other medicinal products

Potent inducers of CYP3A4 may reduce the efficacy of erlotinib whereas potent inhibitors of CYP3A4 may lead to increased toxicity. Concomitant treatment with these types of agents should be avoided (see section 4.5).

#### Other forms of interactions

Erlotinib is characterised by a decrease in solubility at pH above 5. Medicinal products that alter the pH of the upper Gastro-Intestinal (GI) tract, like proton pump inhibitors, H<sub>2</sub> antagonists and antacids, may alter the solubility of erlotinib and hence its bioavailability. Increasing the dose of <Product name> when co-administered with such agents is not likely to compensate for the loss of exposure. Combination of erlotinib with proton pump inhibitors should be avoided. The effects of concomitant administration of erlotinib with H<sub>2</sub> antagonists and antacids are unknown; however, reduced bioavailability is likely. Therefore, concomitant administration of these combinations should be avoided (see section 4.5). If the use of antacids is considered necessary during treatment with <Product name>, they should be taken at least 4 hours before or 2 hours after the daily dose of <Product name>.

#### **The tablets contain lactose.**

Patients with rare hereditary problems of galactose intolerance, Lapp lactase deficiency or glucose-galactose malabsorption should not take this medicine.

### **4.5 Interaction with other medicinal products and other forms of interaction**

Interaction studies have only been performed in adults.

#### Erlotinib and other CYP substrates

Erlotinib is a potent inhibitor of CYP1A1, and a moderate inhibitor of CYP3A4 and CYP2C8, as well as a strong inhibitor of glucuronidation by UGT1A1 *in vitro*.

The physiological relevance of the strong inhibition of CYP1A1 is unknown due to the very limited expression of CYP1A1 in human tissues.

When erlotinib was co-administered with ciprofloxacin, a moderate CYP1A2 inhibitor, the erlotinib exposure [AUC] increased significantly by 39%, while no statistically significant change in C<sub>max</sub> was found. Similarly, the exposure to the active metabolite increased by about 60% and 48% for AUC and C<sub>max</sub>, respectively. The clinical relevance of this increase has not been established. Caution should be exercised when ciprofloxacin or potent CYP1A2 inhibitors (e.g. fluvoxamine) are combined with erlotinib. If adverse reactions related to erlotinib are observed, the dose of erlotinib may be reduced.

Pre-treatment or co-administration of <Product name> did not alter the clearance of the prototypical CYP3A4 substrates, midazolam and erythromycin, but did appear to decrease the oral bioavailability of midazolam by up to 24%. In another clinical study, erlotinib was shown not to affect pharmacokinetics of the concomitantly administered CYP3A4/2C8 substrate paclitaxel. Significant interactions with the clearance of other CYP3A4 substrates are therefore unlikely.

The inhibition of glucuronidation may cause interactions with medicinal products which are substrates of UGT1A1 and exclusively cleared by this pathway. Patients with low expression levels of UGT1A1 or genetic glucuronidation disorders (e.g. Gilbert's disease) may exhibit increased serum concentrations of bilirubin and must be treated with caution.

Erlotinib is metabolised in the liver by the hepatic cytochromes in humans, primarily CYP3A4 and to a lesser extent by CYP1A2. Extrahepatic metabolism by CYP3A4 in intestine, CYP1A1 in lung, and CYP1B1 in tumour tissue also potentially contribute to the metabolic clearance of erlotinib. Potential interactions may occur with active substances which are metabolised by, or are inhibitors or inducers of, these enzymes.

Potent inhibitors of CYP3A4 activity decrease erlotinib metabolism and increase erlotinib plasma concentrations. In a clinical study, the concomitant use of erlotinib with ketoconazole (200 mg orally twice daily for 5 days), a potent CYP3A4 inhibitor, resulted in an increase of erlotinib exposure (86% of AUC and 69% of  $C_{max}$ ). Therefore, caution should be used when erlotinib is combined with a potent CYP3A4 inhibitor, e.g. azole antifungals (i.e. ketoconazole, itraconazole, voriconazole), protease inhibitors, erythromycin or clarithromycin. If necessary the dose of erlotinib should be reduced, particularly if toxicity is observed.

Potent inducers of CYP3A4 activity increase erlotinib metabolism and significantly decrease erlotinib plasma concentrations. In a clinical study, the concomitant use of erlotinib and rifampicin (600 mg orally once daily for 7 days), a potent CYP3A4 inducer, resulted in a 69% decrease in the median erlotinib AUC. Co-administration of rifampicin with a single 450 mg dose of <Product name> resulted in a mean erlotinib exposure (AUC) of 57.5% of that after a single 150 mg <Product name> dose in the absence of rifampicin treatment. Co-administration of <Product name> with CYP3A4 inducers should therefore be avoided. For patients who require concomitant treatment with <Product name> and a potent CYP3A4 inducer such as rifampicin an increase in dose to 300 mg should be considered while their safety (including renal and liver functions and serum electrolytes) is closely monitored, and if well tolerated for more than 2 weeks, further increase to 450 mg could be considered with close safety monitoring. Reduced exposure may also occur with other inducers e.g. phenytoin, carbamazepine, barbiturates or St. John's Wort (*hypericum perforatum*). Caution should be observed when these active substances are combined with erlotinib. Alternate treatments lacking potent CYP3A4 inducing activity should be considered when possible.

#### Erlotinib and coumarin-derived anticoagulants

Interaction with coumarin-derived anticoagulants including warfarin leading to increased International Normalized Ratio (INR) and bleeding events, which in some cases were fatal, have been reported in patients receiving <Product name>. Patients taking coumarin-derived anticoagulants should be monitored regularly for any changes in prothrombin time or INR.

#### Erlotinib and statins

The combination of <Product name> and a statin may increase the potential for statin-induced myopathy, including rhabdomyolysis, which was observed rarely.

#### Erlotinib and smokers

Results of a pharmacokinetic interaction study indicated a significant 2.8-, 1.5- and 9-fold reduced  $AUC_{inf}$ ,  $C_{max}$  and plasma concentration at 24 hours, respectively, after administration of <Product name> in smokers as compared to non-smokers (see section 5.2). Therefore, patients who are still smoking should be encouraged to stop smoking as early as possible before initiation of treatment with <Product name>, as plasma erlotinib concentrations are reduced otherwise. The clinical effect of the decreased exposure has not been formally assessed but it is likely to be clinically significant.

#### Erlotinib and P-glycoprotein inhibitors

Erlotinib is a substrate for the P-glycoprotein active substance transporter. Concomitant administration of inhibitors of Pgp, e.g. cyclosporine and verapamil, may lead to altered distribution and/or altered elimination of erlotinib. The consequences of this interaction for e.g. CNS toxicity have not been established. Caution should be exercised in such situations.

#### Erlotinib and medicinal products altering pH

Erlotinib is characterised by a decrease in solubility at pH above 5. Medicinal products that alter the pH of the upper Gastro-Intestinal (GI) tract may alter the solubility of erlotinib and hence its bioavailability. Co-administration of erlotinib with omeprazole, a proton pump inhibitor (PPI), decreased the erlotinib exposure [AUC] and maximum concentration [ $C_{max}$ ] by 46% and 61%, respectively. There was no change to  $T_{max}$  or half-life. Concomitant administration of <Product name> with 300 mg ranitidine, an H<sub>2</sub>-receptor antagonist, decreased erlotinib exposure [AUC] and maximum concentrations [ $C_{max}$ ] by 33% and 54%, respectively. Increasing the dose of <Product name> when co-administered with such agents is not likely to compensate for this loss of exposure. However, when <Product name> was dosed in a staggered manner 2 hours before or 10 hours after ranitidine 150 mg b.i.d., erlotinib exposure [AUC] and maximum concentrations [ $C_{max}$ ] decreased only by 15% and 17%, respectively. The effect of antacids on the absorption of erlotinib has not been investigated but absorption may be impaired, leading to lower plasma levels. In summary, the combination of erlotinib with proton pump inhibitors should be avoided. If the use of antacids is considered necessary during treatment with <Product name>, they should be taken at least 4 hours before or 2 hours after the daily dose of <Product name>. If the use of ranitidine is considered, it should be used in a staggered manner; i.e. <Product name> must be taken at least 2 hours before or 10 hours after ranitidine dosing.

#### Erlotinib and Gemcitabine

In a Phase Ib study, there were no significant effects of gemcitabine on the pharmacokinetics of erlotinib nor were there significant effects of erlotinib on the pharmacokinetics of gemcitabine.

#### Erlotinib and Carboplatin/Paclitaxel

Erlotinib increases platinum concentrations. In a clinical study, the concomitant use of erlotinib with carboplatin and paclitaxel led to an increase of total platinum  $AUC_{0-48}$  of 10.6%. Although statistically significant, the magnitude of this difference is not considered to be clinically relevant. In clinical practice, there may be other co-factors leading to an increased exposure to carboplatin like renal impairment. There were no significant effects of carboplatin or paclitaxel on the pharmacokinetics of erlotinib.

#### Erlotinib and Capecitabine

Capecitabine may increase erlotinib concentrations. When erlotinib was given in combination with capecitabine, there was a statistically significant increase in erlotinib AUC and a borderline increase in  $C_{max}$  when compared with values observed in another study in which erlotinib was given as single agent. There were no significant effects of erlotinib on the pharmacokinetics of capecitabine.

#### Erlotinib and proteasome inhibitors

Due to the working mechanism, proteasome inhibitors including bortezomib may be expected to influence the effect of EGFR inhibitors including erlotinib. Such influence is supported by limited clinical data and preclinical studies showing EGFR degradation through the proteasome.

#### **4.6 Fertility, pregnancy and lactation**

##### Pregnancy

There are no adequate data for the use of erlotinib in pregnant women. Studies in animals have shown no evidence of teratogenicity or abnormal parturition. However, an adverse effect on the pregnancy can not be excluded as rat and rabbit studies have shown increased embryo/foetal lethality, (see section 5.3). The potential risk for humans is unknown.

##### Women of childbearing potential

Women of childbearing potential must be advised to avoid pregnancy while on <Product name>. Adequate contraceptive methods should be used during therapy, and for at least 2 weeks after completing therapy. Treatment should only be continued in pregnant women if the potential benefit to the mother outweighs the risk to the foetus.

##### Breast-feeding

It is not known whether erlotinib is excreted in human milk. Because of the potential harm to the infant, mothers should be advised against breast-feeding while receiving <Product name>.

##### Fertility

Studies in animals have shown no evidence of impaired fertility. However, an adverse effect on the fertility can not be excluded as animal studies have shown effects on reproductive parameters (see section 5.3). The potential risk for humans is unknown.

#### **4.7 Effects on ability to drive and use machines**

No studies on the effects on the ability to drive and use machines have been performed; however erlotinib is not associated with impairment of mental ability.

#### **4.8 Undesirable effects**

##### Non-small cell lung cancer (<Product name> administered as monotherapy):

In a randomized double-blind study (BR.21; erlotinib administered as second line therapy), rash (75%) and diarrhoea (54%) were the most commonly reported adverse drug reactions (ADRs). Most were Grade 1/2 in severity and manageable without intervention. Grade 3/4 rash and diarrhoea occurred in 9% and 6%, respectively in erlotinib -treated patients and each resulted in study discontinuation in 1% of patients. Dose reduction for rash and diarrhoea was needed in 6% and 1% of patients, respectively. In study BR.21, the median time to onset of rash was 8 days, and the median time to onset of diarrhoea was 12 days.

In general, rash manifests as a mild or moderate erythematous and papulopustular rash, which may occur or worsen in sun exposed areas. For patients who are exposed to sun, protective clothing, and/or use of sun screen (e.g. mineral-containing) may be advisable.

Adverse reactions occurring more frequently ( $\geq 3\%$ ) in erlotinib -treated patients than in the placebo group in the pivotal study BR.21, and in at least 10% of patients in the erlotinib group, are summarised by National Cancer Institute-Common Toxicity Criteria (NCI-CTC) Grade in Table 1.

The following terms are used to rank the undesirable effects by frequency: very common

( $\geq 1/10$ ); common ( $\geq 1/100$  to  $< 1/10$ ); uncommon ( $\geq 1/1,000$  to  $< 1/100$ ); rare ( $\geq 1/10,000$  to  $< 1/1,000$ ); very rare ( $< 1/10,000$ ) including isolated reports.

Within each frequency grouping, adverse reactions are presented in order of decreasing seriousness.

Table 1: Very common ADRs in study BR.21

NCI-CTC Grade	Erlotinib N = 485			Placebo N = 242		
	Any Grade	3	4	Any Grade	3	4
MedDRA Preferred Term	%	%	%	%	%	%
Total patients with any AE	99	40	22	96	36	22
<i>Infections and infestations</i>						
Infection*	24	4	0	15	2	0
<i>Metabolism and nutrition disorders</i>						
Anorexia	52	8	1	38	5	<1
<i>Eye disorders</i>						
Keratoconjunctivitis sicca	12	0	0	3	0	0
Conjunctivitis	12	<1	0	2	<1	0
<i>Respiratory, thoracic and mediastinal disorders</i>						
Dyspnoea	41	17	11	35	15	11
Cough	33	4	0	29	2	0
<i>Gastrointestinal disorders</i>						
Diarrhoea**	54	6	<1	18	<1	0
Nausea	33	3	0	24	2	0
Vomiting	23	2	<1	19	2	0
Stomatitis	17	<1	0	3	0	0
Abdominal pain	11	2	<1	7	1	<1
<i>Skin and subcutaneous tissue disorders</i>						
Rash***	75	8	<1	17	0	0
Pruritus	13	<1	0	5	0	0
Dry skin	12	0	0	4	0	0
<i>General disorders and administration site conditions</i>						
Fatigue	52	14	4	45	16	4

\* Severe infections, with or without neutropenia, have included pneumonia, sepsis, and cellulitis.

\*\* Can lead to dehydration, hypokalemia and renal failure.

\*\*\* Rash included dermatitis acneiform.

In two other double-blind, randomized, placebo-controlled Phase III studies BO18192 (SATURN) and BO25460 (IUNO); erlotinib was administered as maintenance after first-line chemotherapy. These studies were conducted in a total of 1532 patients with advanced, recurrent or metastatic NSCLC following first-line standard platinum- based chemotherapy,

no new safety signals were identified.

The most frequent ADRs seen in patients treated with erlotinib in studies BO18192 and BO25460 were rash and diarrhoea (see Table 2). No Grade 4 rash or diarrhoea was observed in either study. Rash and diarrhoea resulted in discontinuation of erlotinib in 1% and <1% of patients, respectively, in study BO18192, while no patients discontinued for rash or diarrhoea in BO25460. Dose modifications (interruptions or reductions) for rash and diarrhoea were needed in 8.3% and 3% of patients, respectively, in study BO18192 and 5.6% and 2.8% of patients, respectively, in study BO25460.

Table 2: Most frequent ADRs in Studies BO18192 (SATURN) and BO25460 (IUNO)

	BO18192 (SATURN)*		BO25460 (IUNO)*	
	Erlotinib n=433	Placebo n=445	Erlotinib n=322	Placebo n=319
	%	%	%	%
<i>Rash</i> , all grades	49.2	5.8	39.4	10.0
Grade 3	6.0	0	5.0	1.6
<i>Diarrhoea</i> , all grades	20.3	4.5	24.2	4.4
Grade 3	1.8	0	2.5	0.3

\*Safety analysis population

In an open-label, randomized phase III study, ML20650 conducted in 154 patients, the safety of erlotinib for first-line treatment of NSCLC patients with EGFR activating mutations was assessed in 75 patients; no new safety signals were observed in these patients.

The most frequent ADRs seen in patients treated with erlotinib in study ML20650 were rash and diarrhoea (any Grade 80% and 57%, respectively), most were Grade 1/2 in severity and manageable without intervention. Grade 3 rash and diarrhoea occurred in 9% and 4% of patients, respectively. No Grade 4 rash or diarrhoea was observed. Both rash and diarrhoea resulted in discontinuation of erlotinib in 1% of patients. Dose modifications (interruptions or reductions) for rash and diarrhoea were needed in 11% and 7% of patients, respectively.

Pancreatic cancer (<Product name> administered concurrently with gemcitabine):

The most common adverse reactions in pivotal study PA.3 in pancreatic cancer patients receiving erlotinib 100 mg plus gemcitabine were fatigue, rash and diarrhoea. In the erlotinib plus gemcitabine arm, Grade 3/4 rash and diarrhoea were each reported in 5% of patients. The median time to onset of rash and diarrhoea was 10 days and 15 days, respectively. Rash and diarrhoea each resulted in dose reductions in 2% of patients, and resulted in study discontinuation in up to 1% of patients receiving erlotinib plus gemcitabine.

Adverse reactions occurring more frequently ( $\geq 3\%$ ) in erlotinib 100 mg plus gemcitabine-treated patients than in the placebo plus gemcitabine group in the pivotal study PA.3, and in at least 10% of patients in the erlotinib 100 mg plus gemcitabine group, are summarised by National Cancer Institute- Common Toxicity Criteria (NCI-CTC) Grade in Table 3.

The following terms are used to rank the undesirable effects by frequency: very common ( $\geq 1/10$ ); common ( $\geq 1/100$  to  $< 1/10$ ); uncommon ( $\geq 1/1,000$  to  $< 1/100$ ); rare ( $\geq 1/10,000$  to  $< 1/1,000$ ); very rare ( $< 1/10,000$ ) including isolated reports.

Within each frequency grouping, adverse reactions are presented in order of decreasing seriousness.

Table 3: Very common ADRs in study PA.3 (100 mg cohort)

NCI-CTC Grade	Erlotinib N = 259			Placebo N = 256		
	Any Grade	3	4	Any Grade	3	4
<b>MedDRA Preferred Term</b>	<b>%</b>	<b>%</b>	<b>%</b>	<b>%</b>	<b>%</b>	<b>%</b>
Total patients with any AE	99	48	22	97	48	16
<i>Infections and infestations</i>						
Infection*	31	3	<1	24	6	<1
<i>Metabolism and nutrition disorders</i>						
Weight decreased	39	2	0	29	<1	0
<i>Psychiatric disorders</i>						
Depression	19	2	0	14	<1	0
<i>Nervous system disorders</i>						
Neuropathy	13	1	<1	10	<1	0
Headache	15	<1	0	10	0	0
<i>Respiratory, thoracic and mediastinal disorders</i>						
Cough	16	0	0	11	0	0
<i>Gastrointestinal disorders</i>						
Diarrhoea**	48	5	<1	36	2	0
Stomatitis	22	<1	0	12	0	0
Dyspepsia	17	<1	0	13	<1	0
Flatulence	13	0	0	9	<1	0
<i>Skin and subcutaneous tissue disorders</i>						
Rash***	69	5	0	30	1	0
Alopecia	14	0	0	11	0	0
<i>General disorders and administration site conditions</i>						
Fatigue	73	14	2	70	13	2
Pyrexia	36	3	0	30	4	0
Rigors	12	0	0	9	0	0

\* Severe infections, with or without neutropenia, have included pneumonia, sepsis, and cellulitis.

\*\* Can lead to dehydration, hypokalemia and renal failure.

\*\*\* Rash included dermatitis acneiform.

#### Other Observations:

Safety evaluation of erlotinib is based on the data from more than 1500 patients treated with at least one 150 mg dose of erlotinib monotherapy and more than 300 patients who received erlotinib 100 or 150 mg in combination with gemcitabine.

The following adverse reactions have been observed in patients who received erlotinib

administered as single agent and patients who received erlotinib concurrently with chemotherapy.

Very common ADRs from the BR 21 and PA 3 studies are presented in Tables 1 and 3, other ADRs including those from other studies are summarized in Table 4.

Within each frequency grouping, adverse reactions are presented in order of decreasing seriousness

Table 4: Summary of ADRs per frequency category:

<b>Body System</b>	<b>Very common (≥1/10)</b>	<b>Common (≥1/100 to &lt;1/10)</b>	<b>Uncommon (≥1/1,000 to &lt;1/100)</b>	<b>Rare (≥1/10,000 to &lt;1/1,000)</b>	<b>Very rare (&lt;1/10,000)</b>
Eye disorders		-Keratitis -Conjunctivitis <sup>1</sup>	-Eyelash changes <sup>2</sup>		-Corneal perforations -Corneal ulcerations -Uveitis
Respiratory, thoracic and mediastinal disorders		-Epistaxis	- Interstitial lung disease (ILD) <sup>3</sup>		
Gastro-intestinal disorders	-Diarrhoea <sup>7</sup>	-Gastro-intestinal bleeding <sup>4, 7</sup>	-Gastro-intestinal perforations <sup>7</sup>		
Hepato biliary disorders	-Liver function test abnormalities <sup>5</sup>			-Hepatic failure <sup>6</sup>	
Skin and subcutaneous tissue disorders		-Alopecia -Dry skin <sup>1</sup> -Paronychia -Folliculitis -Acne/ Dermatitis acneiform -Skin fissures	-Hirsutism -Eyebrow changes -Brittle and Loose nails -Mild skin reactions such as hyperpigmentation	-Palmar plantar erythrodysesthesia syndrome	-Stevens-Johnson syndrome/Toxic epidermal necrolysis <sup>7</sup>
Renal and urinary disorders		-Renal insufficiency <sup>1</sup>	-Nephritis <sup>1</sup> -Proteinuria <sup>1</sup>		

<sup>1</sup> In clinical study PA.3.

<sup>2</sup> Including in-growing eyelashes, excessive growth and thickening of the eyelashes.

<sup>3</sup> Including fatalities, in patients receiving erlotinib for treatment of NSCLC or other advanced solid tumours (see section 4.4). A higher incidence has been observed in patients in Japan (see section 4.4).

<sup>4</sup> In clinical studies, some cases have been associated with concomitant warfarin administration and some with concomitant NSAID administration (see section 4.5).

<sup>5</sup> Including increased alanine aminotransferase [ALT], aspartate aminotransferase [AST] and bilirubin. These were very common in clinical study PA.3 and common in clinical study BR.21. Cases were mainly mild to moderate in severity, transient in nature or associated with liver metastases.

<sup>6</sup> Including fatalities. Confounding factors included pre-existing liver disease or concomitant hepatotoxic medications (see section 4.4).

<sup>7</sup> Including fatalities (see section 4.4).

#### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the national reporting system listed in Appendix V.

## **4.9 Overdose**

### Symptoms

Single oral doses of <Product name> up to 1000 mg erlotinib in healthy subjects, and up to 1600 mg in cancer patients have been tolerated. Repeated twice daily doses of 200 mg in healthy subjects were poorly tolerated after only a few days of dosing. Based on the data from these studies, severe adverse reactions such as diarrhoea, rash and possibly increased activity of liver aminotransferases may occur above the recommended dose.

### Management

In case of suspected overdose, <Product name> should be withheld and symptomatic treatment initiated.

## **5. PHARMACOLOGICAL PROPERTIES**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: antineoplastic agent protein kinase inhibitor, ATC code: L01XE03

#### Mechanism of action

Erlotinib is an epidermal growth factor receptor/human epidermal growth factor receptor type 1 (EGFR also known as HER1) tyrosine kinase inhibitor. Erlotinib potently inhibits the intracellular phosphorylation of EGFR. EGFR is expressed on the cell surface of normal cells and cancer cells. In non-clinical models, inhibition of EGFR phosphotyrosine results in cell stasis and/or death.

EGFR mutations may lead to constitutive activation of anti-apoptotic and proliferation signaling pathways. The potent effectiveness of erlotinib in blocking EGFR-mediated signalling in these EGFR mutation positive tumours is attributed to the tight binding of erlotinib to the ATP-binding site in the mutated kinase domain of the EGFR. Due to the blocking of downstream-signaling, the proliferation of cells is stopped, and cell death is induced through the intrinsic apoptotic pathway. Tumour regression is observed in mouse models of enforced expression of these EGFR activating mutations.

#### Clinical efficacy

- First-line Non-Small Cell Lung Cancer (NSCLC) therapy for patients with EGFR activating mutations (<Product name> administered as monotherapy):

The efficacy of erlotinib in first-line treatment of patients with EGFR activating mutations in NSCLC was demonstrated in a phase III, randomized, open-label trial (ML20650, EURTAC). This study was conducted in Caucasian patients with metastatic or locally advanced NSCLC (stage IIIB and IV) who have not received previous chemotherapy or any systemic antitumour

therapy for their advanced disease and who present mutations in the tyrosine kinase domain of the EGFR (exon 19 deletion or exon 21 mutation). Patients were randomized 1:1 to receive erlotinib 150 mg daily or up to 4 cycles of platinum based doublet chemotherapy. The primary endpoint was investigator assessed PFS. The efficacy results are summarized in Table 5.

Figure 1: Kaplan-Meier curve for investigator assessed PFS in trial ML20650 (EURTAC) (April 2012 cut-off)




Table 5: Efficacy results of erlotinib versus chemotherapy in trial ML20650 (EURTAC)

		<b>Erlotinib</b>	<b>Chemotherapy</b>	<b>Hazard Ratio (95% CI)</b>	<b>p-value</b>
Pre-planned Interim Analysis (35% OS maturity) (n=153)  Cut-off date: Aug 2010		n=77	n=76		
	Primary endpoint: Progression Free Survival (PFS, median in months)* Investigator Assessed**	9.4	5.2	0.42 [0.27-0.64]	p<0.0001
	Independent Review **	10.4	5.4	0.47 [0.27-0.78]	p=0.003
	Best Overall Response Rate (CR/PR)	54.5%	10.5%		p<0.0001
	Overall Survival (OS) (months)	22.9	18.8	0.80 [0.47-1.37]	p=0.4170
Exploratory Analysis (40%)		n=86	n=87		

OS maturity) (n=173)  Cut-off date: Jan 2011	PFS (median in months), Investigator assessed	9.7	5.2	0.37 [0.27-0.54]	p<0.0001
	Best Overall Response Rate (CR/PR)	58.1%	14.9%		p<0.0001
	OS (months)	19.3	19.5	1.04 [0.65-1.68]	p=0.8702
Updated Analysis (62% OS maturity) (n=173)  Cut-off date: April 2012		n=86	n=87		
	PFS (median in months)	10.4	5.1	0.34 [0.23-0.49]	p<0.0001
	OS*** (months)	22.9	20.8	0.93 [0.64-1.36]	p=0.7149

CR=complete response; PR=partial response

\* A 58% reduction in the risk of disease progression or death was observed

\*\* Overall concordance rate between investigator and IRC assessment was 70%

\*\*\* A high crossover was observed with 82% of the patients in the chemotherapy arm receiving subsequent therapy with an EGFR tyrosine kinase inhibitor and all but 2 of those patients had subsequent erlotinib.

- Maintenance NSCLC therapy after first-line chemotherapy (<Product name> administered as monotherapy):

The efficacy and safety of erlotinib as maintenance after first-line chemotherapy for NSCLC was investigated in a randomized, double-blind, placebo-controlled trial (BO18192, SATURN). This study was conducted in 889 patients with locally advanced or metastatic NSCLC who did not progress after 4 cycles of platinum-based doublet chemotherapy. Patients were randomized 1:1 to receive erlotinib 150 mg or placebo orally once daily until disease progression. The primary endpoint of the study included progression free survival (PFS) in all patients. Baseline demographic and disease characteristics were well balanced between the two treatment arms. Patients with ECOG PS>1, significant hepatic or renal comorbidities were not included in the study.

In this study, the overall population showed a benefit for the primary PFS end point (HR= 0.71 p<0.0001) and the secondary OS end-point (HR= 0.81 p= 0.0088). However the largest benefit was observed in a predefined exploratory analysis in patients with EGFR activating mutations (n= 49) demonstrating a substantial PFS benefit (HR=0.10, 95% CI, 0.04 to 0.25; p<0.0001 and an overall survival HR of 0.83 (95% CI, 0.34 to 2.02). 67% of placebo patients in the EGFR mutation positive subgroup received second or further line treatment with EGFR-TKIs.

The BO25460 (IUNO) study was conducted in 643 patients with advanced NSCLC whose tumors did not harbor an EGFR-activating mutation (exon 19 deletion or exon 21 L858R mutation) and who had not experienced disease progression after four cycles of platinum-based chemotherapy.

The objective of the study was to compare the overall survival of first line maintenance therapy with erlotinib versus erlotinib administered at the time of disease progression. The study did not meet its primary endpoint. OS of erlotinib in first line maintenance was not superior to erlotinib as second line treatment in patients whose tumor did not harbor an

EGFR-activating mutation (HR= 1.02, 95% CI, 0.85 to 1.22, p=0.82). The secondary endpoint of PFS showed no difference between erlotinib and placebo in maintenance treatment (HR=0.94, 95 % CI, 0.80 to 1.11; p=0.48).

Based on the data from the BO25460 (IUNO) study, erlotinib use is not recommended for first-line maintenance treatment in patients without an EGFR activating mutation.

- NSCLC treatment after failure of at least one prior chemotherapy regimen (<Product name> administered as monotherapy):

The efficacy and safety of erlotinib as second/third-line therapy was demonstrated in a randomised, double-blind, placebo-controlled trial (BR.21), in 731 patients with locally advanced or metastatic NSCLC after failure of at least one chemotherapy regimen. Patients were randomised 2:1 to receive erlotinib 150 mg or placebo orally once daily. Study endpoints included overall survival, progression-free survival (PFS), response rate, duration of response, time to deterioration of lung cancer-related symptoms (cough, dyspnoea and pain), and safety. The primary endpoint was survival.

Demographic characteristics were well balanced between the two treatment groups. About two-thirds of the patients were male and approximately one-third had a baseline ECOG performance status (PS) of 2, and 9% had a baseline ECOG PS of 3. Ninety-three percent and 92% of all patients in the erlotinib and placebo groups, respectively, had received a prior platinum-containing regimen and 36% and 37% of all patients, respectively, had received a prior taxane therapy.

The adjusted hazard ratio (HR) for death in the erlotinib group relative to the placebo group was 0.73 (95% CI, 0.60 to 0.87) (p = 0.001). The percent of patients alive at 12 months was 31.2% and 21.5%, for the erlotinib and placebo groups, respectively. The median overall survival was 6.7 months in the erlotinib group (95% CI, 5.5 to 7.8 months) compared with 4.7 months in the placebo group (95% CI, 4.1 to 6.3 months).

The effect on overall survival was explored across different patient subsets. The effect of erlotinib on overall survival was similar in patients with a baseline performance status (ECOG) of 2-3 (HR = 0.77, 95% CI 0.6-1.0) or 0-1 (HR = 0.73, 95% CI 0.6-0.9), male (HR = 0.76, 95% CI 0.6-0.9) or female patients (HR = 0.80, 95% CI 0.6-1.1), patients < 65 years of age (HR = 0.75, 95% CI 0.6-0.9) or older patients (HR = 0.79, 95% CI 0.6-1.0), patients with one prior regimen (HR = 0.76, 95% CI 0.6-1.0) or more than one prior regimen (HR = 0.75, 95% CI 0.6-1.0), Caucasian (HR = 0.79, 95% CI 0.6-1.0) or Asian patients (HR = 0.61, 95% CI 0.4-1.0), patients with adenocarcinoma (HR = 0.71, 95% CI 0.6-0.9) or squamous cell carcinoma (HR = 0.67, 95% CI 0.5-0.9), but not in patients with other histologies (HR 1.04, 95% CI 0.7-1.5), patients with stage IV disease at diagnosis (HR = 0.92, 95% CI 0.7-1.2) or < stage IV disease at diagnosis (HR = 0.65, 95% CI 0.5-0.8). Patients who never smoked had a much greater benefit from erlotinib (survival HR = 0.42, 95% CI 0.28-0.64) compared with current or ex-smokers (HR = 0.87, 95% CI 0.71-1.05).

In the 45% of patients with known EGFR-expression status, the hazard ratio was 0.68 (95% CI 0.49-0.94) for patients with EGFR-positive tumours and 0.93 (95% CI 0.63-1.36) for patients with EGFR-negative tumours (defined by IHC using EGFR pharmDx kit and defining EGFR-negative as less than 10% tumour cells staining). In the remaining 55% of patients with unknown EGFR-expression status, the hazard ratio was 0.77 (95% CI 0.61-0.98).

The median PFS was 9.7 weeks in the erlotinib group (95% CI, 8.4 to 12.4 weeks) compared with 8.0 weeks in the placebo group (95% CI, 7.9 to 8.1 weeks).

The objective response rate by RECIST in the erlotinib group was 8.9% (95% CI, 6.4 to 12.0). The first 330 patients were centrally assessed (response rate 6.2%); 401 patients were investigator- assessed (response rate 11.2%).

The median duration of response was 34.3 weeks, ranging from 9.7 to 57.6+ weeks. The proportion of patients who experienced complete response, partial response or stable disease was 44.0% and 27.5%, respectively, for the erlotinib and placebo groups (p = 0.004).

A survival benefit of erlotinib was also observed in patients who did not achieve an objective tumour response (by RECIST). This was evidenced by a hazard ratio for death of 0.82 (95% CI, 0.68 to 0.99) among patients whose best response was stable disease or progressive disease.

Erlotinib resulted in symptom benefits by significantly prolonging time to deterioration in cough, dyspnoea and pain, versus placebo.

-Pancreatic cancer (<Product name> administered concurrently with gemcitabine in study PA.3):

The efficacy and safety of erlotinib in combination with gemcitabine as a first-line treatment was assessed in a randomised, double-blind, placebo-controlled trial in patients with locally advanced, unresectable or metastatic pancreatic cancer. Patients were randomised to receive erlotinib or placebo once daily on a continuous schedule plus gemcitabine IV (1000 mg/m<sup>2</sup>, Cycle 1 - Days 1, 8, 15, 22, 29, 36 and 43 of an 8 week cycle; Cycle 2 and subsequent cycles - Days 1, 8 and 15 of a 4 week cycle [approved dose and schedule for pancreatic cancer, see the gemcitabine SPC]). Erlotinib or placebo was taken orally once daily until disease progression or unacceptable toxicity. The primary endpoint was overall survival.


Baseline demographic and disease characteristics of the patients were similar between the 2 treatment groups, 100 mg erlotinib plus gemcitabine or placebo plus gemcitabine, except for a slightly larger proportion of females in the erlotinib/gemcitabine arm compared with the placebo/gemcitabine arm:


<b>Baseline</b>	<b>Erlotinib</b>	<b>Placebo</b>
Females	51%	44%
Baseline ECOG performance status (PS) = 0	31%	32%
Baseline ECOG performance status (PS) = 1	51%	51%
Baseline ECOG performance status (PS) = 2	17%	17%
Metastatic disease at baseline	77%	76%

Survival was evaluated in the intent-to-treat population based on follow-up survival data. Results are shown in the table below (results for the group of metastatic and locally advanced patients are derived from exploratory subgroup analysis).

<b>Outcome</b>	<b>Erlotinib (months)</b>	<b>Placebo (months)</b>	<b>Δ (months)</b>	<b>CI of Δ</b>	<b>HR</b>	<b>CI of HR</b>	<b>P-value</b>
<b>Overall Population</b>							
Median overall survival	6.4	6.0	0.41	-0.54-1.64	0.82	0.69-0.98	0.028
Mean overall survival	8.8	7.6	1.16	-0.05-2.34			

Metastatic Population							
Median overall survival	5.9	5.1	0.87	-0.26-1.56	0.80	0.66-0.98	0.029
Mean overall survival	8.1	6.7	1.43	0.17-2.66			
Locally Advanced Population							
Median overall survival	8.5	8.2	0.36	-2.43-2.96	0.93	0.65-1.35	0.713
Mean overall survival	10.7	10.5	0.19	-2.43-2.69			





In a post-hoc analysis, patients with favourable clinical status at baseline (low pain intensity, good QoL and good PS) may derive more benefit from erlotinib. The benefit is mostly driven by the presence of a low pain intensity score.

In a post-hoc analysis, patients on erlotinib who developed a rash had a longer overall survival compared to patients who did not develop rash (median OS 7.2 months vs 5 months, HR: 0.61). 90% of patients on erlotinib developed rash within the first 44 days. The median time to onset of rash was 10 days.

#### Paediatric population

The European Medicines Agency has waived the obligation to submit the results of studies with erlotinib in all subsets of the paediatric population in Non Small Cell Lung Cancer and Pancreatic cancer indications (see section 4.2 for information on paediatric use).

## **5.2 Pharmacokinetic properties**

Absorption: After oral administration, erlotinib peak plasma levels are obtained in approximately 4 hours after oral dosing. A study in normal healthy volunteers provided an estimate of the absolute bioavailability of 59%. The exposure after an oral dose may be increased by food.

Distribution: Erlotinib has a mean apparent volume of distribution of 232 l and distributes into tumour tissue of humans. In a study of 4 patients (3 with non-small cell lung cancer [NSCLC], and 1 with laryngeal cancer) receiving 150 mg daily oral doses of erlotinib, tumour samples from surgical excisions on Day 9 of treatment revealed tumour concentrations of erlotinib that averaged 1185 ng/g of tissue. This corresponded to an overall average of 63% (range 5-161%) of the steady state observed peak plasma concentrations. The primary active metabolites were present in tumour at concentrations averaging 160 ng/g tissue, which corresponded to an overall average of 113% (range 88-130%) of the observed steady state peak plasma concentrations. Plasma protein binding is approximately 95%. Erlotinib binds to serum albumin and alpha-1 acid glycoprotein (AAG).

Biotransformation: Erlotinib is metabolised in the liver by the hepatic cytochromes in humans, primarily CYP3A4 and to a lesser extent by CYP1A2. Extrahepatic metabolism by CYP3A4 in intestine, CYP1A1 in lung, and 1B1 in tumour tissue potentially contribute to the metabolic clearance of erlotinib.

There are three main metabolic pathways identified: 1) O-demethylation of either side chain or both, followed by oxidation to the carboxylic acids; 2) oxidation of the acetylene moiety followed by hydrolysis to the aryl carboxylic acid; and 3) aromatic hydroxylation of the phenyl-acetylene moiety. The primary metabolites OSI-420 and OSI-413 of erlotinib produced by O-demethylation of either side chain have comparable potency to erlotinib in non-clinical *in vitro* assays and *in vivo* tumour models. They are present in plasma at levels that are <10% of erlotinib and display similar pharmacokinetics as erlotinib.

Elimination: Erlotinib is excreted predominantly as metabolites via the faeces (>90%) with renal elimination accounting for only a small amount (approximately 9%) of an oral dose. Less than 2% of the orally administered dose is excreted as parent substance. A population pharmacokinetic analysis in 591 patients receiving single agent erlotinib shows a mean apparent clearance of 4.47 l/hour with a median half-life of 36.2 hours. Therefore, the time to reach steady state plasma concentration would be expected to occur in approximately 7-8 days.

#### Pharmacokinetics in special populations:

Based on population pharmacokinetic analysis, no clinically significant relationship between predicted apparent clearance and patient age, bodyweight, gender and ethnicity were observed. Patient factors, which correlated with erlotinib pharmacokinetics, were serum total bilirubin, AAG and current smoking. Increased serum concentrations of total bilirubin and AAG concentrations were associated with a reduced erlotinib clearance. The clinical relevance of these differences is unclear. However, smokers had an increased rate of erlotinib clearance. This was confirmed in a pharmacokinetic study in non-smoking and currently cigarette smoking healthy subjects receiving a single oral dose of 150 mg erlotinib. The geometric mean of the  $C_{max}$  was 1056 ng/mL in the non-smokers and 689 ng/mL in the smokers with a mean ratio for smokers to non-smokers of 65.2% (95% CI: 44.3 to 95.9,  $p = 0.031$ ). The geometric mean of the  $AUC_{0-inf}$  was 18726 ng•h/mL in the non-smokers and 6718 ng•h/mL in the smokers with a mean ratio of 35.9% (95% CI: 23.7 to 54.3,  $p < 0.0001$ ). The geometric mean of the  $C_{24h}$  was 288 ng/mL in the non-smokers and 34.8 ng/mL in the smokers with a mean ratio of 12.1% (95% CI: 4.82 to 30.2,  $p = 0.0001$ ).

In the pivotal Phase III NSCLC trial, current smokers achieved erlotinib steady state trough plasma concentration of 0.65 µg/mL (n=16) which was approximately 2-fold less than the former smokers or patients who had never smoked (1.28 µg/mL, n=108). This effect was accompanied by a 24% increase in apparent erlotinib plasma clearance. In a phase I dose escalation study in NSCLC patients who were current smokers, pharmacokinetic analyses at steady-state indicated a dose proportional increase in erlotinib exposure when the erlotinib dose was increased from 150 mg to the maximum tolerated dose of 300 mg. Steady-state trough plasma concentrations at a 300 mg dose in current smokers in this study was 1.22 µg/mL (n=17).

Based on the results of pharmacokinetic studies, current smokers should be advised to stop smoking while taking erlotinib, as plasma concentrations could be reduced otherwise.

Based on population pharmacokinetic analysis, the presence of an opioid appeared to increase exposure by about 11%.

A second population pharmacokinetic analysis was conducted that incorporated erlotinib data from 204 pancreatic cancer patients who received erlotinib plus gemcitabine. This analysis

demonstrated that covariants affecting erlotinib clearance in patients from the pancreatic study were very similar to those seen in the prior single agent pharmacokinetic analysis. No new covariate effects were identified. Co-administration of gemcitabine had no effect on erlotinib plasma clearance.

Paediatric population: There have been no specific studies in paediatric patients.

Elderly population: There have been no specific studies in elderly patients.

Hepatic impairment: Erlotinib is primarily cleared by the liver. In patients with solid tumours and with moderately impaired hepatic function (Child-Pugh score 7-9), geometric mean erlotinib AUC<sub>0-t</sub> and C<sub>max</sub> was 27000 ng•h/mL and 805 ng/mL, respectively, as compared to 29300 ng•h/mL and 1090 ng/mL in patients with adequate hepatic function including patients with primary liver cancer or hepatic metastases. Although the C<sub>max</sub> was statistically significant lower in moderately hepatic impaired patients, this difference is not considered clinically relevant. No data are available regarding the influence of severe hepatic dysfunction on the pharmacokinetics of erlotinib. In population pharmacokinetic analysis, increased serum concentrations of total bilirubin were associated with a slower rate of erlotinib clearance.

Renal impairment: Erlotinib and its metabolites are not significantly excreted by the kidney, as less than 9% of a single dose is excreted in the urine. In population pharmacokinetic analysis, no clinically significant relationship was observed between erlotinib clearance and creatinine clearance, but there are no data available for patients with creatinine clearance <15 ml/min.

### **5.3 Preclinical safety data**

Chronic dosing effects observed in at least one animal species or study included effects on the cornea (atrophy, ulceration), skin (follicular degeneration and inflammation, redness, and alopecia), ovary (atrophy), liver (liver necrosis), kidney (renal papillary necrosis and tubular dilatation), and gastrointestinal tract (delayed gastric emptying and diarrhoea). Red blood cell parameters were decreased and white blood cells, primarily neutrophils, were increased. There were treatment-related increases in ALT, AST and bilirubin. These findings were observed at exposures well below clinically relevant exposures.

Based on the mode of action, erlotinib has the potential to be a teratogen. Data from reproductive toxicology tests in rats and rabbits at doses near the maximum tolerated dose and/or maternally toxic doses showed reproductive (embryotoxicity in rats, embryo resorption and foetotoxicity in rabbits) and developmental (decrease in pup growth and survival in rats) toxicity, but was not teratogenic and did not impair fertility. These findings were observed at clinically relevant exposures.

Erlotinib tested negative in conventional genotoxicity studies. Two-year carcinogenicity studies with erlotinib conducted in rats and mice were negative up to exposures exceeding human therapeutic exposure (up to 2-fold and 10-fold higher, respectively, based on C<sub>max</sub> and/or AUC).

A mild phototoxic skin reaction was observed in rats after UV irradiation.

## **6. PHARMACEUTICAL PARTICULARS**

### **6.1 List of excipients**

*Tablet core:*

Lactose Monohydrate  
Cellulose, Microcrystalline (E460)  
Sodium Starch Glycolate Type A  
Magnesium Stearate (E470b)

*Tablet coat:*

Poly(vinyl alcohol) (E1203)  
Titanium Dioxide (E171)  
Macrogol 3350 (E1521)  
Talc (E553b)  
Methacrylic Acid – Ethyl Acrylate Copolymer (1:1), Type A  
Sodium Hydrogen Carbonate

## **6.2 Incompatibilities**

Not applicable.

## **6.3 Shelf life**

3 years

## **6.4 Special precautions for storage**

This medicinal product does not require any special storage conditions.

## **6.5 Nature and contents of container**

HU/H/0461/001-004/DC:

Aluminium - OPA/Alu/PVC blisters of 30 tablets, packed into carton boxes.  
Aluminium - OPA/Alu/PVC perforated unit dose blisters of 30 x1 tablets, packed into carton boxes.

HU/H/0462/001-004/DC:

Aluminium - OPA/Alu/PVC blisters of 30, 90 and 100 tablets, packed into carton boxes.

HU/H/0471/001, 003, 004/DC:

Aluminium - OPA/Alu/PVC blisters of 30 tablets, packed into carton boxes.

HU/H/0472/001, 003, 004/DC:

Aluminium - OPA/Alu/PVC blisters of 20, 30, 40, 60 and 90 tablets, packed into carton boxes.

HU/H/0476/001, 003, 004/DC:

Aluminium - OPA/Alu/PVC blisters of 30 tablets, packed into carton boxes.

HU/H/0496/003-004/DC:

Aluminium - OPA/Alu/PVC blisters of 30 tablets, packed into carton boxes.

Not all pack sizes may be marketed.

## **6.6 Special precautions for disposal**

No special requirements for disposal.

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

**7. MARKETING AUTHORISATION HOLDER**

<To be completed nationally>

**8. MARKETING AUTHORISATION NUMBER(S)**

<To be completed nationally>

**9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE  
AUTHORISATION**

<To be completed nationally>

**10. DATE OF REVISION OF THE TEXT**

<To be completed nationally>

## Package leaflet: Information for the user

<Product name> 25 mg film-coated tablets  
<Product name> 50 mg film-coated tablets  
<Product name> 100 mg film-coated tablets  
<Product name> 150 mg film-coated tablets

Erlotinib

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

### What is in this leaflet

1. What <Product name> is and what it is used for
2. What you need to know before you take <Product name>
3. How to take <Product name>
4. Possible side effects
5. How to store <Product name>
6. Contents of the pack and other information

#### 1. What <Product name> is and what it is used for

<Product name> contains the active substance erlotinib. <Product name> is a medicine used to treat cancer by preventing the activity of a protein called epidermal growth factor receptor (EGFR). This protein is known to be involved in the growth and spread of cancer cells.

<Product name> is indicated for adults. This medicine can be prescribed to you if you have non-small cell lung cancer at an advanced stage. It can be prescribed as initial therapy or as therapy if your disease remains largely unchanged after initial chemotherapy, provided your cancer cells have specific EGFR mutations. It can also be prescribed if previous chemotherapy has not helped to stop your disease.

This medicine can also be prescribed to you in combination with another treatment called gemcitabine if you have cancer of the pancreas at a metastatic stage.

#### 2. What you need to know before you take <Product name>

##### Do not take <Product name>

- if you are allergic to erlotinib or any of the ingredients of this medicine (listed in section 6).

##### Warnings and precautions

- if you are taking other medicines that may increase or decrease the amount of erlotinib in your blood or influence its effect (for example antifungals like ketoconazole, protease inhibitors, erythromycin, clarithromycin, phenytoin, carbamazepine, barbiturates, rifampicin, ciprofloxacin, omeprazole, ranitidine, St. John's Wort or proteasome inhibitors), talk to your doctor. In some cases these

medicines may reduce the efficacy or increase the side effects of <Product name> and your doctor may need to adjust your treatment. Your doctor might avoid treating you with these medicines while you are receiving <Product name>.

- if you are taking anticoagulants (a medicine which helps to prevent thrombosis or blood clotting e.g. warfarin), <Product name> may increase your tendency to bleed. Talk to your doctor, he will need to regularly monitor you with some blood tests.
- if you are taking statins (medicines to lower your blood cholesterol), <Product name> may increase the risk of statin related muscle problems, which on rare occasions can lead to serious muscle breakdown (rhabdomyolysis) resulting in kidney damage, talk to your doctor.
- if you use contact lenses and/or have a history of eye problems such as severe dry eyes, inflammation of the front part of the eye (cornea) or ulcers involving the front part of the eye, tell your doctor.

See also below “Other medicines and <Product name>”.

You should tell your doctor:

- if you have sudden difficulty in breathing associated with cough or fever because your doctor may need to treat you with other medicines and interrupt your <Product name> treatment;
- if you have diarrhoea because your doctor may need to treat you with anti-diarrhoeal (for example loperamide);
- immediately, if you have severe or persistent diarrhoea, nausea, loss of appetite, or vomiting because your doctor may need to interrupt your <Product name> treatment and may need to treat you in the hospital;
- if you have severe pain in the abdomen, severe blistering or peeling of skin. Your doctor may need to interrupt or stop your treatment;
- if you develop acute or worsening redness and pain in the eye, increased eye watering, blurred vision and/or sensitivity to light, please tell your doctor or nurse immediately as you may need urgent treatment (see Possible Side Effects below).
- if you are also taking a statin and experience unexplained muscle pain, tenderness, weakness or cramps. Your doctor may need to interrupt or stop your treatment.

See also section 4 “Possible side effects”.

#### Liver or kidney disease

It is not known whether <Product name> has a different effect if your liver or kidneys are not functioning normally. The treatment with this medicine is not recommended if you have a severe liver disease or severe kidney disease.

#### Glucuronidation disorder like Gilbert’s syndrome

Your doctor must treat you with caution if you have a glucuronidation disorder like Gilbert’s syndrome.

#### Smoking

You are advised to stop smoking if you are treated with <Product name> as smoking could decrease the amount of your medicine in the blood.

#### **Children and adolescents**

<Product name> has not been studied in patients under the age of 18 years. The treatment with this medicine is not recommended for children and adolescents.

#### **Other medicines and <Product name>**

Tell your doctor or pharmacist if you are taking, have recently taken any other medicines or might take any other medicines.

**<Product name> with food and drink**

Do not take <Product name> with food. See also section 3 'How to take <Product name>'

**Pregnancy and breast-feeding**

Avoid pregnancy while being treated with <Product name>. If you could become pregnant, use adequate contraception during treatment, and for at least 2 weeks after taking the last tablet.

If you become pregnant while you are being treated with <Product name>, immediately inform your doctor who will decide if the treatment should be continued.

Do not breast-feed if you are being treated with <Product name>.

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

**Driving and using machines**

<Product name> has not been studied for its possible effects on the ability to drive and use machines but it is very unlikely that your treatment will affect this ability.

**<Product name> contains a sugar called lactose monohydrate.**

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

**3. How to take <Product name>**

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist if you are not sure.

The tablet should be taken at least one hour before or two hours after the ingestion of food.

The recommended dose is one tablet of <Product name> 150 mg each day if you have non-small cell lung cancer.

The recommended dose is one tablet of <Product name> 100 mg each day if you have metastatic pancreatic cancer. <Product name> is given in combination with gemcitabine treatment.

Your doctor may adjust your dose in 50 mg steps.

For the different dose regimens <Product name> is available in strengths of 25 mg, 100 mg or 150 mg.

For MSs where 25 mg, 50 mg and 100 mg are not registered:

<Product name> is not available in 25 mg, 50 mg and 100 mg strengths. For these dosages, you should take other medicinal products available on the market. Ask your doctor or pharmacist.

For oral use.

**If you take more <Product name> than you should**

Contact your doctor or pharmacist immediately.

You may have increased side effects and your doctor may interrupt your treatment.

**If you forget to take <Product name>**

If you miss one or more doses of <Product name>, contact your doctor or pharmacist as soon as possible. Do not take a double dose to make up for a forgotten dose.

**If you stop taking <Product name>**

It is important to keep taking <Product name> every day, as long as your doctor prescribes it for you.

If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

**4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Contact your doctor as soon as possible if you suffer from any of the below side effects. In some cases your doctor may need to reduce your dose of <Product name> or interrupt treatment:

- Diarrhoea and vomiting (very common: may affect more than 1 out of 10 people). Persistent and severe diarrhoea may lead to low blood potassium and impairment of your kidney function, particularly if you receive other chemotherapy treatments at the same time. If you experience more severe or persistent diarrhoea **contact your doctor immediately** as your doctor may need to treat you in the hospital.
- Eye irritation due to conjunctivitis/keratoconjunctivitis (very common: may affect more than 1 out of 10 people) and keratitis (common: may affect up to 1 in 10 people).
- Form of lung irritation called interstitial lung disease (uncommon in European patients; common in Japanese patients: may affect up to 1 in 100 people in Europe and up to 1 in 10 in Japan). This disease can also be linked to the natural progression of your medical condition and can have a fatal outcome in some cases. If you develop symptoms such as sudden difficulty in breathing associated with cough or fever contact your doctor immediately as you could suffer from this disease. Your doctor may decide to permanently stop your treatment with <Product name>.
- Gastrointestinal perforations have been observed (uncommon: may affect up to 1 in 100 people). Tell your doctor if you have severe pain in your abdomen. Also, tell your doctor if you had peptic ulcers or diverticular disease in the past, as this may increase this risk.
- In rare cases liver failure was observed (rare: may affect up to 1 in 1,000 people). If your blood tests indicate severe changes in your liver function, your doctor may need to interrupt your treatment.

**Very common side effects (may affect more than 1 in 10 people):**

- Rash which may occur or worsen in sun exposed areas. If you are exposed to sun, protective clothing, and/or use of sun screen (e.g. mineral-containing) may be advisable
- Infection
- Loss of appetite, decreased weight
- Depression
- Headache, altered skin sensation or numbness in the extremities
- Difficulty in breathing, cough
- Nausea
- Mouth irritation
- Stomach pain, indigestion and flatulence

- Abnormal blood tests for the liver function
- Itching, dry skin and loss of hair
- Tiredness, fever, rigors

**Common side effects (may affect up to 1 in 10 people):**

- Bleeding from the nose
- Bleeding from the stomach or the intestines
- Inflammatory reactions around the fingernail
- Infection of hair follicles
- Acne
- Cracked skin (skin fissures)
- Reduced kidney function (when given outside the approved indications in combination with chemotherapy)

**Uncommon side effects (may affect up to 1 in 100 people):**

- Eyelash changes
- Excess body and facial hair of a male distribution pattern
- Eyebrow changes
- Brittle and loose nails

**Rare side effects (may affect up to 1 in 1,000 people):**

- Flushed or painful palms or soles (Palmar plantar erythrodysesthesia syndrome)

**Very rare side effects (may affect up to 1 in 10,000 people):**

- Cases of perforation or ulceration of the cornea
- Severe blistering or peeling of skin (suggestive of Stevens-Johnson syndrome)
- Inflammation of the coloured part of the eye

**Reporting of side effects**

If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the national reporting system listed in Appendix V. By reporting side effects you can help provide more information on the safety of this medicine.

**5. How to store <Product name>**

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton and blister after EXP. The expiry date refers to the last day of that month.

This medicine does not require any special storage conditions.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

**6. Contents of the pack and other information**

**What <Product name> contains**

- The active substance is erlotinib.

<Product name> 25 mg film-coated tablets

Each film-coated tablet contains 25 mg of erlotinib (as erlotinib hydrochloride).

<Product name> 50 mg film-coated tablets

Each film-coated tablet contains 50 mg of erlotinib (as erlotinib hydrochloride).

<Product name> 100 mg film-coated tablets

Each film-coated tablet contains 100 mg of erlotinib (as erlotinib hydrochloride).

<Product name> 150 mg film-coated tablets

Each film-coated tablet contains 150 mg of erlotinib (as erlotinib hydrochloride).

- The other ingredients are:

*Tablet core:* lactose monohydrate, cellulose microcrystalline (E460), sodium starch glycolate type A, magnesium Stearate (E470b).

*Tablet coat:* poly(vinyl alcohol) (E1203), titanium dioxide (E171), macrogol 3350 (E1521), talc (E553b), methacrylic acid – ethyl acrylate copolymer (1:1), type A, sodium hydrogen carbonate

### **What <Product name> looks like and contents of the pack**

<Product name> 25 mg film-coated tablets: White to yellowish, round biconvex, film-coated tablet, engraved with “25” on one side. The diameter of the tablet is 6.1 mm ± 5 %.

<Product name> 50 mg film-coated tablets: White to yellowish, round biconvex, film-coated tablet, engraved with “50” on one side. The diameter of the tablet is 7.6 mm ± 5 %.

<Product name> 100 mg film-coated tablets: White to yellowish, round biconvex, film-coated tablet, engraved with “100” on one side. The diameter of the tablet is 8.9 mm ± 5 %.

<Product name> 150 mg film-coated tablets: White to yellowish, round biconvex, film-coated tablet, engraved with “150” on one side. The diameter of the tablet is 10.5 mm ± 5 %.

#### HU/H/0461/001-004/DC:

The tablets are available in Aluminium - OPA/Alu/PVC blisters of 30 tablets, packed into carton boxes.

The tablets are available in Aluminium - OPA/Alu/PVC perforated unit dose blisters of 30 x1 tablets, packed into carton boxes.

#### HU/H/0462/001-004/DC:

The tablets are available in Aluminium - OPA/Alu/PVC blisters of 30, 90 and 100 tablets, packed into carton boxes.

#### HU/H/0471/001, 003, 004/DC:

The tablets are available in Aluminium - OPA/Alu/PVC blisters of 30 tablets, packed into carton boxes.

#### HU/H/0472/001, 003, 004/DC:

The tablets are available in Aluminium - OPA/Alu/PVC blisters of 20, 30, 40, 60 and 90 tablets, packed into carton boxes.

#### HU/H/0476/001, 003, 004/DC:

The tablets are available in Aluminium - OPA/Alu/PVC blisters of 30 tablets, packed into carton boxes.

#### HU/H/0496/003-004/DC:

The tablets are available in Aluminium - OPA/Alu/PVC blisters of 30 tablets, packed into carton boxes.

Not all pack sizes may be marketed.

### **Marketing Authorisation Holder and Manufacturer**

<To be completed nationally>

**<This medicinal product is authorised in the Member States of the EEA under the following names:>**

<To be completed nationally>

**This leaflet was last revised in {MM/YYYY}.**

**PARTICULARS TO APPEAR ON THE OUTER PACKAGING**

**OUTER CARTON**

**1. NAME OF THE MEDICINAL PRODUCT**

<Product name> 25 mg film-coated tablets  
<Product name> 50 mg film-coated tablets  
<Product name> 100 mg film-coated tablets  
<Product name> 150 mg film-coated tablets  
Erlotinib

**2. STATEMENT OF ACTIVE SUBSTANCE(S)**

Each film-coated tablet contains 25 mg erlotinib (as erlotinib hydrochloride).  
Each film-coated tablet contains 50 mg erlotinib (as erlotinib hydrochloride).  
Each film-coated tablet contains 100 mg erlotinib (as erlotinib hydrochloride).  
Each film-coated tablet contains 150 mg erlotinib (as erlotinib hydrochloride).

**3. LIST OF EXCIPIENTS**

Contains lactose monohydrate.  
See leaflet for further information.

**4. PHARMACEUTICAL FORM AND CONTENTS**

HU/H/0461/001-004/DC:  
30 and 30 x1 film-coated tablets

HU/H/0462/001-004/DC:  
30, 90 and 100 film-coated tablets

HU/H/0471/001, 003, 004/DC:  
30 film-coated tablets

HU/H/0472/001, 003, 004/DC:  
20, 30, 40, 60 and 90 film-coated tablets

HU/H/0476/001, 003, 004/DC:  
30 film-coated tablets

HU/H/0496/003-004/DC:  
30 film-coated tablets

**5. METHOD AND ROUTE(S) OF ADMINISTRATION**

For oral use.

Read the package leaflet before use.

**6. SPECIAL WARNING THAT THE MEDICINAL PRODUCT MUST BE STORED OUT OF THE SIGHT AND REACH OF CHILDREN**

Keep out of the sight and reach of children.

**7. OTHER SPECIAL WARNING(S), IF NECESSARY**

**8. EXPIRY DATE**

EXP

**9. SPECIAL STORAGE CONDITIONS**

**10. SPECIAL PRECAUTIONS FOR DISPOSAL OF UNUSED MEDICINAL PRODUCTS OR WASTE MATERIALS DERIVED FROM SUCH MEDICINAL PRODUCTS, IF APPROPRIATE**

**11. NAME AND ADDRESS OF THE MARKETING AUTHORISATION HOLDER**

<To be completed nationally>

**12. MARKETING AUTHORISATION NUMBER(S)**

<To be completed nationally>

**13. BATCH NUMBER**

Lot

**14. GENERAL CLASSIFICATION FOR SUPPLY**

Medicinal product subject to medical prescription

**15. INSTRUCTIONS ON USE**

**16. INFORMATION IN BRAILLE**

<Product name> 25 mg

<Product name> 50 mg  
<Product name> 100 mg  
<Product name> 150 mg

**17. UNIQUE IDENTIFIER – 2D BARCODE**

<2D barcode carrying the unique identifier included.>

**18. UNIQUE IDENTIFIER - HUMAN READABLE DATA**

PC: {number}  
SN: {number}  
NN: {number}

**MINIMUM PARTICULARS TO APPEAR ON BLISTERS OR STRIPS**

**BLISTERS**

**1. NAME OF THE MEDICINAL PRODUCT**

<Product name> 25 mg film-coated tablets  
<Product name> 50 mg film-coated tablets  
<Product name> 100 mg film-coated tablets  
<Product name> 150 mg film-coated tablets  
Erlotinib

**2. NAME OF THE MARKETING AUTHORISATION HOLDER**

<To be completed nationally>

**3. EXPIRY DATE**

EXP

**4. BATCH NUMBER**

Lot

**5. OTHER**